

Haematologica  
HAEMATOL/2017/174177  
Version 4

A distinct subtype of Epstein Barr virus positive T/NK-cell lymphoproliferative disorder: Adult patients with chronic active Epstein Barr virus infection-like features

Keisuke Kawamoto, Hiroaki Miyoshi, Takaharu Suzuki, Yasuji Kozai,  
Koji Kato, Masaharu Miyahara, Toshiaki Yujiri, Ilseung Choi,  
Katsumichi Fujimaki, Tsuyoshi Muta, Masaaki Kume, Sayaka  
Moriguchi, Shinobu Tamura, Takeharu Kato, Hiroyuki Tagawa, Junya  
Makiyama, Yuji Kanisawa, Yuya Sasaki, Daisuke Kurita, Kyohei  
Yamada, Joji Shimono, Hirohito Sone, Jun Takizawa, Masao Seto,  
Hiroshi Kimura, and Koichi Ohshima

Disclosures: 1. This work was partly supported by The Tsukada Medical Foundation and The Yasuda Medical Foundation. 2.3. None

Contributions: Contribution: K.K., H.M., and K.O. were responsible for conception and design; K.K., T.S., Y.K., M.M., T.Y., N.O., I.C., K.F., T.M., T.N., M.K., S.M., S.T., T.K., H.T., J.M., Y.K., Y.S., D.K., K.Y., and J.S. performed collection and assembly of data; K.K. and H.M. performed data analysis and interpretation; K.K. wrote the manuscript; J.T., H.S, H.K., M.S. and K.O. supervised the manuscript; and all authors approved the final manuscript.