

Haematologica  
HAEMATOL/2017/173781  
Version 4

C/EBP $\gamma$  is dispensable for steady-state and emergency granulopoiesis

Miroslava Kardosova, Polina Zjablovskaja, Petr Danek, Pavla Angelisova, Lorena Lobo de Figueiredo-Pontes, Robert S. Welner, Tomas Brdicka, Sanghoon Lee, Daniel G. Tenen, and Meritxell Alberich-Jorda

Disclosures: The authors would like to thank the Flow Cytometry Core Facility, IMG AS CR, Prague, Czech Republic for their support with the flow cytometry and sorting presented herein. We would like to thank to Dr. P. Johnson and Dr. J. Salotti for advice in EMSA assays and Dr. V. Korinek for his technical support. The study was supported by a GACR grant (18-08577S) and institutional funding from the IMG AS CR (RVO 68378050) to MA-J, a FAPESP grant (2015/21866-1) to LLF-P, and by a GA UK fellowship (project No. 341015) from Charles University in Prague to MK. DGT was supported by a STaR Investigator Award, an RCE Core grant, an Tier 3 RNA Biology Center grant MOE2014-T3-1-006 from the NRF and MOE, Singapore; and NIH/NCI grant P01 CA66996. This work used instruments provided by C4Sys infrastructure and it was also supported by the Czech Centre for Phenogenomics (CCP, project no. LM2015040), and OP RDI CZ.1.05/2.1.00/19.0395 (Higher quality and capacity for transgenic models).

Contributions: MA-J and DGT conceived and designed the study; MK, PD, PZ, PA, LP, RW, and SL carried out experiments; MK, PA, TB, DGT, and MA-J performed data analysis and interpretation; MK, and MA-J wrote the manuscript.