

Haematologica
HAEMATOL/2018/191809
Version 4

Impact of FLT3-ITD length on prognosis of acute myeloid leukemia

Song-Bai Liu, Hao-Jie Dong, Xie-Bing Bao, Qiao-Cheng Qiu, Hong-Zhi Li, Hong-Jie Shen, Zi-Xuan Ding, Chao Wang, Xiao-Ling Chu, Jing-Qiu Yu, Tao Tao, Zheng Li, Xiao-Wen Tang, Su-Ning Chen, De-Pei Wu, Ling Li, and Sheng-Li Xue

Disclosures: This work was supported by grants from the National Natural Science Foundation of China (Grant No. 81470296, 31701198), Jiangsu Provincial Medical Youth Talent Program (Grant No. QNRC2016719, QNRC2016771), Natural Science Foundation of Jiangsu Province of China (Grant No. BK20170386), a C class sponsored project of Jiangsu provincial Six Talent Peaks (Grant No. 2016-WSN-123), and the China National Clinical Key Subject Project. This work was supported in part by the National Institutes of Health under award number R00CA184411, the Margaret E. Early Medical Research Trust Award, V Scholar Research Award, the Stop Cancer Research Career Development Award and the Gehr Family Center for Leukemia Research (to L.L.). Research reported here includes work performed in the Analytical Cytometry Core at City of Hope supported by the National Cancer Institute of the National Institutes of Health under award number P30CA33572. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We also thank Lynn Smith for the help in manuscript editing and proofreading.

Contributions: S.L., H.D., X.B., Q.Q., designed and performed research, analyzed data, and wrote the manuscript; H.L., H.S., Z.D., C.W., X.C., J.Y., T.T., L.Z., X.W., S.C., performed research, analyzed data, and reviewed the manuscript; L.L. designed the research, analyzed and interpreted data, and wrote the manuscript; S.X. and D.W. analyzed and interpreted data, and wrote the manuscript.