



## editorial

### MILD HYPERHOMOCYSTEINEMIA AND ARTERIAL OCCLUSIVE DISEASE

JACOB SELHUB

USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA, USA

The initial link between homocysteine and vascular disease was made by McCully approximately 25 years ago.<sup>1</sup> He observed that an infant who died as a result of a rare genetic condition of abnormal cobalamin metabolism with homocystinuria exhibited widespread, severe arteriosclerosis analogous to the lesions seen in cases of homocystinuria caused by a genetic cystathionine  $\beta$ -synthase deficiency. Because hyperhomocysteinemia was the only condition common to these two metabolic disorders, McCully proposed that hyperhomocysteinemia resulted in arteriosclerotic disease. Although McCully's hypothesis did not gain immediate support, the association between plasma homocysteine concentration and arteriosclerosis has more recently become the subject of a number of clinical studies. In 1976 Wilcken and Wilcken showed that the concentration of homocysteine-cysteine mixed disulphide after a methionine load was slightly higher in CHD patients than in respective age-sex matched controls.<sup>2</sup> This pioneering work has led to many studies which have been the subject of two important review articles. The first by Ueland *et al.*<sup>3</sup> summarized 17 studies that presented fasting homocysteine concentrations for approximately 1500 patients with various forms of vascular disease and a similar number of normal controls. Fasting homocysteine concentrations were consistently elevated among patients with all types of vascular disease and averaged 31% greater than concentrations among controls. Abnormal homocysteine metabolism was also measured in some studies by presenting individuals with a large oral dose of methionine, which is a homocysteine precursor. Frequencies of abnormal homocysteine response to methionine loading were higher (24%) in patients with vascular disease than in healthy controls (2%). The more recent metanalysis by Boushey *et al.*<sup>4</sup> of 27 studies including prospective and population-based case-control studies concluded that elevations of total plasma homocysteine (tHcy) were considered an independent graded risk factor for arteriosclerotic

vascular disease, with odds ratios for a 5  $\mu\text{mol/L}$  increase in plasma tHcy that were equal to 1.5-1.8 for men and women with CHD, cerebrovascular or peripheral vascular diseases.

Our interest in homocysteine was prompted by the possibility that plasma homocysteine may serve as an indicator of the status and perhaps the intake of a number of vitamins, including folic acid, vitamin B<sub>12</sub> and vitamin B<sub>6</sub>. This possibility derived from the large number of studies which implied that methionine metabolism is tightly regulated,<sup>3,6,7</sup> and from other studies which showed that deficiencies in the above vitamins are often associated with hyperhomocysteinemia.<sup>8-18</sup>

The aims of our studies were therefore: 1) to determine the relationship between plasma homocysteine concentrations and status and intake of folate, vitamin B<sub>12</sub> and vitamin B<sub>6</sub>; 2) to determine the prevalence of hyperhomocysteinemia; 3) to determine the contribution of vitamin status and intake to the prevalence of hyperhomocysteinemia and, 4) to describe the associations between plasma homocysteine, vitamin status and intake and prevalence of carotid artery stenosis.

In our recent studies<sup>19,20</sup> we examined members of the original Framingham Heart Study cohort, a population-based sample of 5209 men and women originally examined in 1948-1952<sup>21</sup> and followed prospectively to the present to assess the occurrence of vascular disease. This study was based on 1401 survivors of the original cohort who participated in the 20<sup>th</sup> biennial examination (1989-90). Homocysteine and carotid ultrasound measures were available for 1041 individuals (418 men and 623 women), aged 67 to 96 years old at the time of data collection. The main findings of these studies were the following.

#### *Homocysteine distribution and prevalence of high homocysteine concentrations*

The mean homocysteine concentration for all subjects was 11.9  $\mu\text{mol/L}$  (median = 11.6  $\mu\text{mol/L}$ ).

Correspondence: Jacob Selhub, PhD, USDA Human Nutrition Research Center on Aging at Tufts University, 711 Washington Street, Boston, MA 02111, USA. Tel. international +1.617.556-3191. Fax international +1.617.556-3344.

Acknowledgments: support for these studies was provided by US Department of Agriculture National Research Initiative Competitive Grants Program No. 92-37200-7582; US Department of Agriculture contract No. 53-3k06-5-10.

Values ranged from 3.5 to 66.9  $\mu\text{mol/L}$ . Homocysteine concentration was higher in men than in women and increased with age. The increase with age remained highly significant ( $p < 0.001$ ) for men and women after adjustment for plasma vitamin concentrations, but the difference between men and women was no longer statistically significant. We defined high homocysteine as concentrations greater than the 90<sup>th</sup> percentile among subjects with all plasma vitamin levels  $>70^{\text{th}}$  percentile (14.0  $\mu\text{mol/L}$ ). Prevalence of high homocysteine was 29.3% for the entire cohort and over 40% for individuals aged 80 years and older.

#### Mean homocysteine concentration by vitamin status and intake

**Folate.** Mean plasma homocysteine concentrations for subjects in the two lowest deciles of plasma folate (below 4.8 nmol/L) were 15.6 and 13.7  $\mu\text{mol/L}$ . These were significantly greater than the mean for subjects in the highest decile, which was 11.0  $\mu\text{mol/L}$  ( $p < 0.01$ ). Mean homocysteine concentrations for subjects in the three lowest deciles of folate intake (less than 253  $\mu\text{g/day}$ ) were 13.7, 12.9 and 13.2  $\mu\text{mol/L}$ , respectively, and were significantly greater than the mean for subjects in the highest intake decile, which was 10.4  $\mu\text{mol/L}$  ( $p < 0.01$ ).

**Vitamin B<sub>12</sub>.** Mean homocysteine concentrations were significantly elevated for subjects in the lowest decile for vitamin B<sub>12</sub> relative to subjects in the highest decile ( $p < 0.01$ ). Mean homocysteine concentrations were 15.4 and 10.9  $\mu\text{mol/L}$  for subjects in the lowest and highest vitamin B<sub>12</sub> deciles. Subjects in the lowest vitamin B<sub>12</sub> decile had vitamin B<sub>12</sub> concentrations below 139 pmol/L. Vitamin B<sub>12</sub> intake appeared to be unrelated to mean homocysteine concentration, even though subjects in the fifth decile had significantly higher homocysteine concentrations than subjects in the highest decile ( $p < 0.05$ ).

**Vitamin B<sub>6</sub>.** Mean homocysteine concentrations were significantly elevated for subjects in the lowest decile for PLP relative to subjects in the highest decile for this vitamin ( $p < 0.01$ ). Mean homocysteine concentrations were 14.3 and 10.9  $\mu\text{mol/L}$  for subjects in the lowest and highest PLP deciles. Subjects in the lowest decile had PLP concentrations below 18.1 nmol/L. For vitamin B<sub>6</sub> intake, mean homocysteine concentrations were significantly elevated in the lowest two deciles ( $p < 0.01$ ) and the third decile ( $p < 0.05$ ). Mean homocysteine concentrations were 13.4, 12.4 and 12.3  $\mu\text{mol/L}$  for subjects in the lowest three deciles; the mean in the highest decile was 10.1  $\mu\text{mol/L}$ . Subjects in the lowest three intake deciles reported consuming less than 1.75 mg/day.

#### Homocysteine concentrations by overall vitamin status

Mean homocysteine and the prevalence of high homocysteine increased dramatically across cate-

gories of the B vitamin index (Table 1). Mean homocysteine concentration was 75% greater in the lowest relative to the highest index categories. The prevalence of high homocysteine was almost 6-fold greater among subjects in the lowest index category compared with subjects in the highest category for plasma index. Sixty-seven percent of the cases of high homocysteine in this cohort of older subjects were associated with at least one vitamin concentration below the 70<sup>th</sup> percentile. Although the prevalence of high homocysteine was substantially greater in lower vitamin categories (4 and 5) than in the middle category, this latter category contributed the largest share of cases of high homocysteine for the index because it included the largest proportion of the cohort.

#### Relationship between plasma homocysteine and prevalence of extracranial stenosis

The prevalence of extracranial carotid stenosis  $\geq 25$  percent was approximately 43 percent and 34 percent in men and women, respectively. In men, the prevalence of stenosis  $\geq 25$  percent was 27 percent (95 percent confidence interval: 17 to 38 percent) in the lowest homocysteine quartile and 58 percent (95 percent confidence interval: 49 to 67 percent) in the highest quartiles ( $p_{\text{trend}} < 0.001$ ). The relationship in women was not as striking as that in men; prevalence of stenosis  $\geq 25$  percent ranged from 31 percent (95 percent confidence interval: 24 to 38 percent) to 39 percent (95 percent confidence interval: 31 to 47 percent) across homocysteine quartiles ( $p_{\text{trend}} = 0.03$ ). While the risk of stenosis appeared to increase in the second homocysteine quartile (9.1 to 11.3  $\mu\text{mol/L}$ ) among men, it did not appear to increase until the third homocysteine quartile (11.4 to 14.3  $\mu\text{mol/L}$ ) among women. Although the prevalence of stenosis appeared somewhat greater among men than women in the upper quartiles of homocysteine, a test of interaction between sex and homocysteine

Table 1. Elevated homocysteine concentrations by B vitamin status.

| B vitamin index | n.  | Mean homocysteine ( $\mu\text{mol/L}$ ) | Prevalence (%) | Prevalence rate ratio | Attributable percent | Population attributable percent |
|-----------------|-----|---|----------------|-----------------------|----------------------|---------------------------------|
| Highest 1       | 89  | 9.4                                     | 10.1           | 1                     | 0.0                  | 0.0                             |
| 2               | 128 | 9.8                                     | 12.5           | 1.2                   | 19.2                 | 1.0                             |
| 3               | 534 | 11.9*                                   | 28.7*          | 2.8                   | 64.8                 | 33.7                            |
| 4               | 144 | 14.9*                                   | 52.1*          | 5.2                   | 80.6                 | 20.6                            |
| Lowest 5        | 70  | 16.5*                                   | 58.6*          | 5.8                   | 52.8                 | 11.6                            |
|                 |     |   |                | (Total)               | 66.9                 |                                 |

1 = index combines plasma folate, vitamin B<sub>12</sub> and PLP concentration; high 1) = all three B vitamins  $>70^{\text{th}}$  percentile; 2 = all vitamins  $>50^{\text{th}}$ , at least 1  $<70^{\text{th}}$  percentile; 3 = vitamins above and below the 50<sup>th</sup> percentile; 4 = all vitamins  $<50^{\text{th}}$  percentile, at least 1  $>30^{\text{th}}$  percentile; low (5) = all three vitamins  $<30^{\text{th}}$  percentile. \*Significantly different from category 1 ( $p < 0.01$ ).

Table 2. Odds ratios of maximal extracranial carotid artery stenosis > 25 percent according to quartile of plasma vitamin concentrations.

|                                 | N     | Adjusted for multiple risk factors |         |         | Adjusted for multiple risk factors plus homocysteine |         |         |
|---------------------------------|-------|------------------------------------|---------|---------|--|---------|---------|
|                                 |       | Odds Ratio                         | 95% CI* | p value | Odds Ratio   | 95% CI* | p value |
| Folate (ug/L)                   |       |                                    |         |         |  |         |         |
| <2.51                           | 1,027 | 1.9                                | 1.3,2.7 | 0.001   | 1.5  | 1.0,2.3 | 0.04    |
| 2.51-4.31                       |       | 1.4                                | 1.0,2.0 | 0.08    | 1.3  | 0.9,1.9 | 0.24    |
| 4.32-7.92                       |       | 1.2                                | 0.8,1.8 | 0.28    | 1.2  | 0.8,1.8 | 0.35    |
| 7.93                            |       | 1.0                                |         |         | 1.0  |         |         |
| P <sub>trend</sub>              |       |                                    |         |         |  | <0.001  | 0.05    |
| Vitamin B <sub>12</sub> (ng/L)  |       |                                    |         |         |  |         |         |
| <290                            | 881   | 1.4                                | 0.9,2.1 | 0.11    | 1.2  | 0.8,1.8 | 0.41    |
| 290-405                         |       | 1.4                                | 0.9,2.0 | 0.14    | 1.2  | 0.8,1.8 | 0.36    |
| 406-572                         |       | 1.3                                | 0.9,2.0 | 0.16    | 1.3  | 0.9,1.9 | 0.24    |
| 573                             |       | 1.0                                |         |         | 1.0  |         |         |
| P <sub>trend</sub>              |       |                                    |         |         |  | 0.11    | 0.47    |
| Vitamin B <sub>6</sub> (nmol/L) |       |                                    |         |         |  |         |         |
| <31.61                          | 967   | 1.6                                | 1.1,2.4 | 0.02    | 1.3  | 0.9,2.0 | 0.15    |
| 31.91-52.19                     |       | 1.1                                | 0.7,1.6 | 0.67    | 1.0  | 0.6,1.4 | 0.80    |
| 52.20-89.80                     |       | 1.2                                | 0.8,1.7 | 0.48    | 1.1  | 0.7,1.6 | 0.71    |
| 89.81                           |       | 1.0                                |         |         | 1.0  |         |         |
| P <sub>trend</sub>              |       |                                    |         |         |  | 0.03    | 0.2     |

\*CI: confidence interval.

indicated that the trends for prevalence of stenosis  $\geq$  25 percent were not significantly different for men and women ( $p=0.07$ ).

The associations between carotid stenosis and plasma vitamins are shown in Table 2. The prevalence of stenosis  $\geq$  25 percent was inversely associated with both folate ( $p_{\text{trend}} < 0.001$ ) and pyridoxal-5'-phosphate ( $p_{\text{trend}} = 0.03$ ) after adjustment for age, sex and other risk factors. The odds ratio for stenosis was 1.9 (95 percent confidence interval: 1.3 to 2.7) in the lowest folate quartile and 1.6 (95 percent confidence interval: 1.1 to 2.4) in the lowest pyridoxal-5'-phosphate quartile. Plasma vitamin B<sub>12</sub> exhibited a weak association with stenosis ( $P_{\text{trend}} = 0.11$ ). The odds ratio for stenosis was 1.4 (95 percent confidence interval: 0.9 to 2.1) in the lowest vitamin B<sub>12</sub> quartile compare with the highest quartile. Adjustment for homocysteine diminished the strength of plasma vitamin associations, but the elevated prevalence of stenosis in the lowest plasma folate quartile remained evident (odds ratio: 1.5; 95 percent confidence interval: 1.0 to 2.3).

These data suggest an important role for nutritional status in homocysteine metabolism. We have demonstrated strong, non-linear, inverse associations between homocysteine concentrations and plasma concentrations of folate, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>. We observed that individuals with low levels of each of these vitamins had high plasma homocysteine concentrations, while those with moderate vitamin levels had dramatically lower homocysteine concentrations. Homocysteine levels did not differ substantially between individuals with moderate and high vitamin concentrations.

The results for folate and vitamin B<sub>6</sub> intake data

are consistent with those for the plasma vitamins. Although it is risky to attribute discrete quantitative values based on this method of dietary assessment,<sup>22</sup> it still may be worth noting that homocysteine concentrations were elevated among individuals with folate intakes up to 280  $\mu\text{g}/\text{day}$ , which is higher than the current RDA of 200 and 180  $\mu\text{g}/\text{day}$  for adult men and women, and vitamin B<sub>6</sub> intakes as high as 1.92 mg/day, which is less than the RDA of 2.0 mg/day for men but greater than the RDA of 1.6 mg/day for women.

Adequate levels of all three vitamins may be needed to obtain an optimal homocysteine concentration. Using the index based on levels of all three vitamins, we estimated that approximately two-thirds of the cases of elevated homocysteine concentration in this cohort were associated with low or moderate plasma levels of one or more of the three vitamins.

Our findings also provide evidence that plasma homocysteine levels are associated with extracranial carotid stenosis in a population-based, elderly cohort. We observed that risk of stenosis  $\geq$  25 percent was increased at homocysteine concentrations previously believed to be normal based on levels of homocysteine among normative samples. As in our previous analysis, we defined elevated plasma homocysteine as concentrations greater than 14  $\mu\text{mol}/\text{L}$  (90<sup>th</sup> percentile among individuals with apparently adequate folate, vitamin B<sub>12</sub> and vitamin B<sub>6</sub> status). Stampfer *et al.*<sup>23</sup> defined elevated homocysteine as concentrations greater than 15.8  $\mu\text{mol}/\text{L}$  (95<sup>th</sup> percentile among non-diseased control subjects). Joosten *et al.*<sup>24</sup> defined elevated homocysteine as concentrations greater than 13.9  $\mu\text{mol}/\text{L}$  (mean plus 2 standard deviations among healthy young controls). Genest and coworkers<sup>25</sup> reported 90<sup>th</sup> and 95<sup>th</sup> percentile values of 15.0 and 19.0  $\mu\text{mol}/\text{L}$  among their normal controls. In the present study we observed that risk of stenosis was elevated at levels of homocysteine between 11.4 and 14.3  $\mu\text{mol}/\text{L}$ . These data will require us to reconsider the current beliefs regarding standards for elevated homocysteine.

We also examined the relationships between specific nutritional determinants of hyperhomocysteinemia and stenosis in this elderly cohort. We further demonstrated that folate and pyridoxal-5'-phosphate were linked to stenosis, in large part, due to their regulation of plasma homocysteine levels as indicated by the diminished odds ratios between stenosis and these vitamins after adjustment for homocysteine levels. Although there was some residual association between plasma folate and stenosis after adjustment for homocysteine, the likelihood ratio test statistic would suggest that addition of folate to a model containing homocysteine did not add any significant contribution. It is likely that measurement error and biological variability in both folate and homocysteine might explain the

residual folate association.

We demonstrated that the majority of these elderly individuals with elevated homocysteine concentrations have insufficient folate, vitamin B<sub>12</sub> or vitamin B<sub>6</sub> status, and others have shown that innocuous vitamin supplementation regimens (including folate, B<sub>12</sub>, and B<sub>6</sub>) effectively lower moderately elevated plasma homocysteine levels to the normal range.<sup>10,18,26-33</sup>

Results of our studies provide the rationale for a randomized, controlled trial of the effect of homocysteine-lowering vitamin therapy on vascular disease morbidity and mortality in hyperhomocysteinemic, elderly individuals. The relevance of this topic was emphasized at the *Meeting on Basic, Laboratory and Clinical Aspects of Thromboembolic Diseases*, held in La Thuile, Italy, on March 17-23, 1996, and two papers presented on that occasion are reported in this issue of *Haematologica*. While the first winter meeting held in 1994 was mainly devoted to genetic disorders responsible for venous thrombosis,<sup>32-36</sup> the 1996 one highlighted the relevance of acquired disorders as a cause of arterial thrombosis.

©1997, Ferrata Storti Foundation

## References

- McCully KS. Vascular pathology of homocysteinemia: implications for the pathogenesis of arteriosclerosis. *Am J Pathol* 1969; 56:111-28.
- Wilcken DEL, Wilcken B. The pathogenesis of coronary artery disease: a possible role for methionine metabolism. *J Clin Invest* 1976; 57:1079-82.
- Ueland PM, Refsum H, Brattstrom L. Plasma homocysteine and cardiovascular disease. In: Francis RB, ed. *Atherosclerotic cardiovascular disease hemostasis and Endothelial Function*. New York:Marcel Dekker; 1992. p. 183-236.
- Boushey J, Beresford SAA, Omenn GS, Motulsky AG. A quantitative assessment of plasma homocysteine as a risk factor for vascular disease. Probable benefits of increasing folic acid intakes. *JAMA* 1995; 274:1049-57.
- Ueland PM, Refsum H. Plasma homocysteine, a risk factor for vascular disease: plasma levels in health, disease, and drug therapy. *J Lab Clin Med* 1989; 114:473-501.
- Mudd SH. Homocystinuria and homocysteine metabolism: selected aspects. In: Nyhan WL, ed. *Heritable disorders of amino acid metabolism*. New York: John Wiley & Sons; 1974. p. 429-51.
- Selhub J, Miller JW. The pathogenesis of homocysteinemia: interruption of the coordinate regulation by S-adenosylmethionine of the remethylation and transsulfuration of homocysteine. *Am J Clin Nutr* 1992; 55:131-8.
- Kang S-S, Wong PWK, Norusis M. Homocysteinemia due to folate deficiency. *Metabolism* 1987; 36:458-62.
- Brattstrom L, Israelsson B, Lindgarde F, Haltberg B. Higher total plasma homocysteine in vitamin B<sub>12</sub> deficiency than in heterozygosity for cystathionine β-synthase deficiency. *Metabolism* 1988; 37:175-8.
- Brattstrom LE, Israelsson B, Jeppsson JO, Hultberg BL. Folic acid-an innocuous means to reduce plasma homocysteine. *Scan J Clin Lab Invest* 1988; 48:215-21.
- Stabler SP, Marcellip D, Podell ER, Allen RH, Savage DG, Lindenbaum J. Elevation of total homocysteine in the serum of patients with cobalamin or folate deficiency detected by capillary gas chromatography-mass spectrometry. *J Clin Invest* 1988; 81:466-74.
- Lindenbaum J, Savage DG, Stabler SP, Allen RH. Diagnosis of cobalamin deficiency: II. Relative sensitivities of serum cobalamin, methylmalonic acid, and total homocysteine concentrations. *Am J Hematol* 1990; 34:99-107.
- Park YK, Linkswiler H. Effect of vitamin B6 depletion in adult man on the excretion of cystathionine and other methionine metabolites. *J Nutrition* 1970; 100:110-6.
- Smolin LA, Benevenga NJ. Accumulation of homocyst(e)ine in vitamin B6 deficiency: a model for the study of cystathionine B-synthase deficiency. *J Nutrition* 1982; 112:1264-72.
- Smolin LA, Benevenga NJ. Factors affecting the accumulation of homocyst(e)ine in rats deficient in vitamin B6. *J Nutrition* 1984; 114:103-11.
- Smolin LA, Benevenga NJ, Berlow S. The use of betaine for the treatment of homocystinuria. *J Pediatr* 1981; 99:467-72.
- Smolin LA, Crenshaw TD, Kurtycz D, Benevenga NJ. Homocyst(e)ine accumulation in pigs fed diets deficient in vitamin B6: relationship to atherosclerosis. *J Nutr* 1983; 113:2122-33.
- Ubbink JB, Vermaak WJH, van der Merwe A, Becker PJ. Vitamin B-12, vitamin B-6 and folate nutritional status in men with hyperhomocysteinemia. *Am J Clin Nutr* 1993; 57:47-53.
- Selhub J, Jacques PF, Wilson PWF, Rush D, Rosenberg IH. Vitamin status and intake as primary determinants of homocysteinemia in an elderly population. *JAMA* 1993; 270:2693-8.
- Selhub J, Jacques PF, Bostom AG, et al. Plasma homocysteine and extracranial carotid stenosis in the framingham heart study. *N Engl J Med* 1995; 332:286-91.
- Dawber TR, Moore FE, Mann GV. Coronary heart disease in the Framingham study. *Am J Public Health* 1957; 47(suppl):4-24.
- Willett W. *Nutritional Epidemiology*. New York:Oxford University Press; 1990. p. 92-126.
- Stampfer MJ, Malinow MR, Willett WC, et al. A prospective study of plasma homocyst(e)ine and risk of myocardial infarction in US physicians. *JAMA* 1992; 268:877-81.
- Joosten E, van den Berg A, Riezler R, et al. Metabolic evidence that deficiencies of vitamin B-12 (cobalamin), folate, and vitamin B-6 occur commonly in elderly people. *Am J Clin Nutr* 1993; 58:468-76.
- Genest JJ, McNamara JR, Salem DN, Wilson PWF, Schaefer EJ, Malinow MR. Plasma homocyst(e)ine levels in men with premature coronary artery disease. *J Am Coll Cardiol* 1990; 16:1114-9.
- Brattstrom L, Israelsson B, Norving B, et al. Impaired homocysteine metabolism in early-onset cerebral and peripheral occlusive arterial disease. Effects of pyridoxine and folic acid treatment. *Atherosclerosis* 1990; 81:51-60.
- Dudman NPB, Wilcken DEL, Wang J, Lynch JF, Macey D, Lundberg P. Disordered methionine/homocysteine metabolism in premature vascular disease: its occurrence, cofactor therapy, and enzymology. *Arterioscler Thromb* 1993; 13:1253-60.
- Franken DG, Boers GHJ, Blom HJ, Trijbels JMF. Effect of various regimens of vitamin B-6 and folic acid on mild hyperhomocysteinemia in vascular patients. *J Inher Metab Dis* 1994; 17:159-62.
- Glueck CJ, Shaw P, Lang JE, Tracy T, Sieve-Smith L, Wang Y. Evidence that homocysteine is an independent risk factor for atherosclerosis in hyperlipidemic patients. *Am J Cardiol* 1995; 55:132-9.
- Lindgren F, Israelsson B, Lindgren A, Hultberg B, Andersson A, Brattstrom L. Plasma homocysteine in acute myocardial infarction: homocysteine-lowering effect of folic acid. *J Intern Med* 1995; 237:381-8.
- Ubbink JB, Vermaak WJH, van der Merwe A, et al. Vitamin requirements for the treatment of hyperhomocysteinemia in humans. *J Nutr* 1994; 124:1927-33.
- Dahlback B. Inherited resistance to activated protein C, a major basis of venous thrombosis, is caused by deficient anticoagulant cofactor function of factor V. *Haematologica* 1995; 80(suppl. to no. 2):102-13.
- Amiral J. Research and development strategies in hemostasis and thrombosis. *Haematologica* 1995; 80:14-24.
- Ascari E, Siragusa S, Piovella F. The epidemiology of deep vein thrombosis and pulmonary embolism. *Haematologica* 1995; 80(suppl. to no. 2):36-41.
- Prandoni P, Villalta S, Polistena P, Bernardi E, Cogo A, Girolami A. Symptomatic deep-vein thrombosis and the post-thrombotic syndrome. *Haematologica* 1995; 80(suppl. to no. 2):42-8.
- Girolami A, Prandoni P, Simioni P, Girolami B, Scarano L, Zanone E. The pathogenesis of venous thromboembolism. *Haematologica* 1995; 80(suppl. to no. 2):25-35.