

Haematologica
HAEMATOL/2018/199190
Version 3

Lysine specific demethylase 1 inactivation enhances differentiation and promotes cytotoxic response when combined with all-trans retinoic acid in acute myeloid leukemia across subtypes

Kimberly N. Smitheman, Tesa M. Severson, Satyajit R. Rajapurkar, Michael T. McCabe, Natalie Karpnich, James Foley, Melissa B. Pappalardi, Ashley Hughes, Wendy Halsey, Elizabeth Thomas, Christopher Traini, Kelly E. Federowicz, Jenny Laraio, Fredrick Mobegi, Geraldine Ferron-Brady, Rabinder K. Prinjha, Christopher L. Carpenter, Ryan G. Kruger, Lodewyk Wessels, and Helai P. Mohammad

Disclosures: none

Contributions: Conception and design: K.N.S., R.G.K., and H.P.M Acquisition of data: K.N.S., N.K., J.F., M.P., A.H., W.H., E.T., C.T., K.E.F., and J.L. Analysis and interpretation of data K.N.S., T.M.S., S.R.R., F.M., G.F.B., L.W., and H.P.M. Writing, review, and/or revision of the manuscript: K.N.S., T.M.S., M.T.M., R.K.P., C.L.C., R.G.K., and H.P.M. Study supervision: C.L.C., R.G.K., and H.P.M.