

Kinase activity is impaired in neutrophils of sepsis patients

Arie J. Hoogendijk,¹ Lonneke A. van Vught,¹ Maryse A. Wiewel,¹ Gwenny M. Fuhler,² Hakima Belkasim-Bohoudi,¹ Janneke Horn,³ Marcus J. Schultz,^{3,4} Brendon P. Scicluna,^{1,5} Maikel P. Peppelenbosch,² Cornelis van 't Veer,¹ Alex F. de Vos¹ and Tom van der Poll^{1,6}

¹Center for Experimental and Molecular Medicine, Academic Medical Center, University of Amsterdam; ²Department of Gastroenterology and Hepatology, Erasmus MC, Erasmus University of Rotterdam; ³Department of Intensive Care, Academic Medical Center, University of Amsterdam; ⁴Laboratory of Experimental Intensive Care and Anesthesiology, Academic Medical Center, University of Amsterdam; ⁵Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Academic Medical Center, University of Amsterdam and ⁶Division of Infectious Diseases, Academic Medical Center, University of Amsterdam, the Netherlands.

Correspondence: TOM VAN DER POLL
t.vanderpoll@amc.uva.nl
doi:10.3324/haematol.2018.201913

Supplementary Table 1:

Kinomics chip kinase substrate proteins (column A) and with RefSeq (protein) or UniProt (column B) identifiers.

Annotated upstream kinases are shown in column C and D using Human Protein Reference Database identifiers.

Columns E-G show significance calls of a substrate (-1 significantly less phosphorylated; 1 significantly more phosphorylated). Cluster assignment is indicated in column H.