

Haematologica
HAEMATOL/2019/216465
Version 4

Increased neonatal level of arginase 2 in cases of childhood acute lymphoblastic leukemia implicates immunosuppression in etiology

Amalie B. Nielsen, Mi Zhou, Adam J. de Smith, Rong Wang, Lucie McCoy, Helen Hansen, Libby Morimoto, Kirsten Grønbæk, Christoffer Johansen, Scott C. Kogan, Catherine Metayer, Paige M. Bracci, Xiaomei Ma, and Joseph L. Wiemels

Disclosures: This work was supported by a grant from the Lundbeck Foundation, Clinical Research Fellowship for A.B.N. Research reported in this publication was supported by funding from the National Institutes of Health (grants R01CA175737, R01ES009137 and P50ES018172) and the Environmental Protection Agency (grant RD83615901) of the United States. The content is solely the responsibility of the authors and does not represent the official views of any funding agencies.

Contributions: A.B.N. performed experiments, analyzed data, and drafted the manuscript with JW; M.Z., A.D. performed experiments and A.D. conceived the study with J.W.; R.W. planned and conducted statistical analysis and co-wrote the sections on statistical analysis and results; H.H. ascertained and managed Guthrie Cards and laboratory activities; L.M., C.J., K.G., S.K., C.M., P.B. contributed to the analysis and reviewed the manuscript; X.M. and J.W. designed the overall study, planned the analysis, and helped to draft the manuscript.