

Impact of corticosteroid pretreatment in pediatric patients with newly diagnosed B-lymphoblastic leukemia: a report from the Children's Oncology Group

Elizabeth A. Raetz,¹ Mignon L. Loh,² Meenakshi Devidas,³ Kelly Maloney,⁴ Leonard A. Mattano Jr.,⁵ Eric Larsen,⁶ Andrew Carroll,⁷ Nyla A. Heerema,⁸ Julie M. Gastier-Foster,⁹ Brent Wood,¹⁰ Michael J. Borowitz,¹¹ Naomi Winick,¹² Stephen P. Hunger¹³ and William L. Carroll¹

¹Department of Pediatrics and Perlmutter Cancer Center, NYU Langone Health, New York, NY; ²Department of Pediatrics, Benioff Children's Hospital, and the Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, CA; ³Department of Biostatistics, Colleges of Medicine, Public Health and Health Professions, University of Florida, Gainesville, FL; ⁴Department of Pediatrics University of Colorado School of Medicine and Children's Hospital Colorado, Aurora, CO; ⁵HARP Pharma Consulting, Mystic, CT; ⁶Department of Pediatrics, Maine Children's Cancer Program, Scarborough, ME; ⁷Department of Genetics, University of Alabama at Birmingham, Birmingham, AL; ⁸Department of Pathology, Wexner Medical Center, Ohio State University, Columbus, OH; ⁹Institute for Genomic Medicine, Nationwide Children's Hospital and Departments of Pathology and Pediatrics, Ohio State University College of Medicine, Columbus, OH; ¹⁰Departments of Pathology and Medicine, University of Washington, Seattle, WA; ¹¹Departments of Pathology and Oncology, Johns Hopkins Medical Institutions, Baltimore, MD; ¹²Departments of Pediatrics, UT Southwestern, Dallas, TX and ¹³Department of Pediatrics and the Center for Childhood Cancer Research, Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, PA, USA

Correspondence: ELIZABETH A. RAETZ - elizabeth.raetz@nyulangone.org
doi:10.3324/haematol.2018.215616

SUPPLEMENTAL DATA

Table S1. Early Treatment Response

	Not Steroid Pretreated N=7542	Steroid Pretreated N=247	P-value
End Induction Remission*			
AALL0331	99.1% (4910/4954)	97.4% (75/77)	0.16
AALL0232	95.2% (2398/2520)	95.8% (158/165)	0.73
Day 29 MRD <0.1%			
AALL0331	92.0% (4547/4943)	87.0% (67/77)	0.11
AALL0232	83.9% (2105/2509)	84.9% (141/166)	0.72
Day 29 MRD <0.01%			
AALL0331	81.6% (4033/4943)	74.0% (57/77)	0.09
AALL0232	73.1% (1833/2509)	70.5% (117/166)	0.47
Rapid Early Response			
AALL0331	87.4% (4319/4944)	85.5% (65/76)	0.63
AALL0232	80.0% (2007/2510)	80.9% (131/162)	0.78

* M1 marrow (<5% blasts) on Day 29.

Table S2. Cox Regression Analysis


Analysis of Maximum Likelihood Estimates				
Factor	P-value	Hazard Ratio	95% Hazard Ratio Confidence Limits	
NCI Risk Group	<0.0001	2.17	1.93	2.45
MRD Day 29 (0.1%)	<0.0001	2.18	1.82	2.60
MRD Day 29 (0.01%)	<0.0001	2.46	2.08	2.91
Steroid Pretreatment	0.70	1.06	0.78	1.44

Figure Legends

Supplemental Figure 1. Outcomes for standard risk patients allocated to a high risk therapy. Five-year event-free (EFS) and overall survival (OS) are shown in corticosteroid pretreated vs. not pretreated patients. A) EFS; B) OS for NCI SR patients assigned to COG AALL0232 due to selected patterns of corticosteroid exposure prior to diagnosis; namely, corticosteroid exposure in the week preceding diagnosis in the absence of a pre-steroid CBC (n=14) and > 48 hours of corticosteroid exposure in weeks -4 to -1 preceding diagnosis (n=31).

Figure S1.

A.



B.

