

Haematologica
HAEMATOL/2018/209650
Version 4

Loss of RAF kinase inhibitor protein is involved in myelomonocytic differentiation and aggravates RAS-driven myeloid leukemogenesis

Veronica Caraffini, Olivia Geiger, Angelika Rosenberger, Stefan Hatzl, Bianca Perfler, Johannes L. Berg, Clarice Lim, Herbert Strobl, Karl Kashofer, Silvia Schauer, Christine Beham-Schmid, Gerald Hoefler, Klaus Geissler, Franz Quehenberger, Walter Kolch, Dimitris Athineos, Karen Blyth, Albert Wölfler, Heinz Sill, and Armin Zebisch

Disclosures: none

Contributions: AZ conceived and supervised the study. VC, OG, AR, BP, JLB, CL, HSt, KK, SS, KB, DA, AW, HSi, and AZ developed the study methodology. VC, OG, AR, SH, BP, JLB, CL, HSt, KK, SS, GH, KG, WK, KB, DA, AW, HSi, and AZ acquired the data. VC, OG, AR, SH, BP, JLB, CL, HSt, KK, CB, GH, WK, KB, DA, AW, HSi, and AZ analysed and interpreted the data. FQ performed statistical analyses. VC and AZ wrote the original draft of the manuscript and all the authors reviewed and edited the manuscript.