

# Post-transplant cyclophosphamide versus anti-thymocyte globulin for graft-versus-host disease prevention in haploidentical transplantation for adult acute lymphoblastic leukemia

Arnon Nagler,<sup>1\*</sup> Abraham S. Kanate,<sup>2\*</sup> Myriam Labopin,<sup>3\*</sup> Fabio Ciceri,<sup>4</sup> Emanuele Angelucci,<sup>5</sup> Yener Koc,<sup>6</sup> Zafer Gülbaz,<sup>7</sup> William Arcese,<sup>8</sup> Johanna Tischer,<sup>9</sup> Pietro Pioltelli,<sup>10</sup> Hakan Ozdogu,<sup>11</sup> Boris Afanasyev,<sup>12</sup> Depei Wu,<sup>13</sup> Mutlu Arat,<sup>14</sup> Zinaida Peric,<sup>15</sup> Sebastian Giebel,<sup>16</sup> Bipin Savani<sup>17</sup> and Mohamad Mohty<sup>18</sup>

<sup>1</sup>Chaim Sheba Medical Center, Tel Aviv University, Tel-Hashomer, Israel; <sup>2</sup>West Virginia University, Morgantown, WV, USA; <sup>3</sup>Department of Hematology and EBMT Paris study office / CEREST-TC, Saint Antoine Hospital, Paris, France; <sup>4</sup>Ospedale San Raffaele s.r.l., Milano, Italy; <sup>5</sup>Ospedale San Martino, Department of Hematology II, Genova, Italy; <sup>6</sup>Medical Park Hospitals, Stem Cell Transplant Unit, Antalya, Turkey; <sup>7</sup>Anadolu Medical Center Hospital, Bone Marrow Transplantation Department, Kocaeli, Turkey; <sup>8</sup>"Tor Vergata" University of Rome, Stem Cell Transplant Unit, Policlinico Universitario Tor Vergata, Rome, Italy; <sup>9</sup>Klinikum Grosshadern, Med. Klinik III, Munich, Germany; <sup>10</sup>Ospedale San Gerardo, Clinica Ematologica dell'Università Milano-Biocca, Monza, Italy; <sup>11</sup>Baskent University Hospital, Hematology Division, Bone Marrow Transplantation Unit, Adana, Turkey; <sup>12</sup>First State Pavlov Medical University of St. Petersburg, Raisa Gorbacheva Memorial Research Institute for Pediatric Oncology, Hematology, and Transplantation, St. Petersburg, Russia; <sup>13</sup>First Affiliated Hospital of Soochow University, Department of Hematology, Suzhou, China; <sup>14</sup>Florence Nightingale Sisli Hospital, Hematopoietic Stem Cell Transplantation Unit, Istanbul, Turkey; <sup>15</sup>University Hospital Center Zagreb, School of Medicine, University of Zagreb, Zagreb, Croatia; <sup>16</sup>Department of Bone Marrow Transplantation and Oncohematology, Maria Skłodowska-Curie Memorial Cancer Center and Institute of Oncology, Gliwice, Poland; <sup>17</sup>Department of Hematology-Oncology, Vanderbilt University Medical Center, Nashville, TN, USA and <sup>18</sup>Department of Hematology and EBMT Paris study office / CEREST-TC, Saint Antoine Hospital, INSERM UMR 938 and Université Pierre et Marie Curie, Paris, France

\*AN, ASK and ML contributed equally as co-first authors.

©2021 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2020.247296

Received: January 12, 2020.

Accepted: April 28, 2020.

Pre-published: April 30, 2020.

Correspondence: ABRAHAM S. KANATE - askanate@hsc.wvu.edu



Male recipient	33.2%[27.6-39]	16.8%[12.5-21.5]	33.2%[26.9-39.7]	12.4%[8.3-17.3]	36.1%[29.8-42.5]	28.3%[22.7-34]	35.6%[29.2-42]	43.7%[37-50.3]	26.7%[20.7-32.7]
Female recipient	26.9%[20-34.3]	7.4%[3.9-12.3]	25.9%[18.4-33.9]	8.1%[4.1-13.9]	35.2%[26.9-43.5]	27.7%[20.3-35.5]	37.2%[28.5-45.8]	43.7%[34.7-52.7]	34.1%[25.7-42.5]
<i>P</i> -value	0.10	0.006	0.19	0.37	0.91	0.72	0.66	0.73	0.18
Male donor	28%[22.2-34]	13.5%[9.4-18.3]	23.9%[17.9-30.4]	7.6%[4.3-12]	34.9%[28.2-41.7]	30.5%[24.2-36.9]	34.6%[27.7-41.5]	38.4%[31.2-45.5]	28.7%[22.1-35.2]
Female donor	34.6%[27.9-41.4]	13.6%[9.2-18.9]	39%[31.1-46.8]	15.5%[10-22]	37%[29.4-44.6]	25.5%[19.3-32.2]	37.4%[29.7-45.2]	49.1%[41.2-57.1]	29.7%[22.3-37]
<i>P</i> -value	0.09	0.97	0.007	0.03	0.95	0.15	0.16	0.049	0.46
No Female (D) → Male (R)	29.5%[24.4-34.8]	12.3%[8.9-16.4]	27.1%[21.6-32.8]	8.7%[5.6-12.8]	35.1%[29.2-41.1]	29.1%[23.8-34.7]	35.7%[29.6-41.8]	40.7%[34.3-47.1]	29.7%[23.8-35.5]
Female (D) → male (R)	34.5%[26.1-43.1]	16%[10-23.1]	38.5%[28.6-48.3]	16.1%[9.4-24.5]	37%[27.5-46.6]	25.5%[17.8-33.9]	37.5%[27.8-47.1]	50.6%[40.8-60.4]	28.6%[19.6-37.7]
<i>P</i> -value	0.18	0.31	0.07	0.07	0.93	0.46	0.58	0.31	0.95
Recipient CMV (-)	25.7%[17.2-35.2]	8.9%[4.1-15.9]	42.2%[30.4-53.5]	7.7%[3.1-15.1]	32%[21.5-42.9]	23.9%[14.9-34]	44.1%[32.3-55.9]	54.4%[43-65.8]	38.8%[27.6-50]
Recipient CMV (+)	32.4%[27.3-37.6]	14.8%[11.1-18.9]	28%[22.6-33.6]	11.7%[8.1-16.1]	37.4%[31.6-43.2]	28.6%[23.6-33.9]	34%[28.2-39.7]	40.8%[34.8-46.9]	26.7%[21.2-32.1]
<i>P</i> -value	0.2	0.16	0.03	0.39	0.45	0.59	0.22	0.15	0.15
Donor CMV (-)	23.9%[16.4-32.4]	10%[5.3-16.4]	31.2%[21.7-41.2]	8%[3.4-15.1]	35.1%[25.4-44.8]	21.7%[14-30.5]	43.3%[32.9-53.6]	50.2%[39.7-60.8]	37.7%[27.7-47.7]
Donor CMV (+)	34.1%[28.7-39.5]	15%[11.2-19.3]	31.4%[25.6-37.4]	12.2%[8.4-16.8]	36.1%[30.1-42.1]	29.9%[24.6-35.4]	34%[28-40]	41.9%[35.6-48.1]	26.6%[20.9-32.2]
<i>P</i> -value	0.03	0.21	0.95	0.41	0.93	0.13	0.21	0.14	0.14

Supplemental Table 1. Univariate analysis of post-transplant outcomes after haploidentical transplantation in ALL patients

Abbreviations: GVHD – graft-versus-host disease; NRM – non-relapse mortality; LFS – leukemia-free survival; OS – overall survival; GRFS – GVHD/relapse-free survival; HCT – hematopoietic cell transplantation; CR – complete remission; ALL – acute lymphoblastic leukemia; KPS – Karnofsky performance score; BM – bone marrow; PB – peripheral blood; MAC – myeloablative conditioning; RIC – reduced intensity conditioning; D – donor; R – recipient; CMV – cytomegalovirus

Supplemental Table 2. Subset analysis of post-transplant outcomes by graft source (Bone marrow vs. peripheral blood)

		100-day Acute GVHD II-IV	2-year Chronic GVHD	2-year Relapse	2-year NRM	2-year LFS	2-year OS	2-year GRFS
<b>Bone Marrow</b>	ATG (n=31)	35.5%[19.1-52.3]	26.7%[11-45.3]	55.4%[34.6-71.9]	20.4%[7.9-37]	24.2%[8-40.3]	34.8%[17.1-52.5]	20.6%[5.2-36]
	PTCy (n=175)	25.4%[19-32.3]	29.6%[22.1-37.6]	33.7%[26.1-41.5]	26.9%[20.1-34.1]	39.4%[31.2-47.6]	48.6%[40.3-57]	32.7%[24.9-40.5]
	<i>P</i> -value	0.17	0.52	<b>0.06</b>	0.41	0.41	0.30	0.75
<b>Peripheral Blood</b>	ATG (n=67)	31.4%[20.3-43]	28.3%[15.9-42]	35.7%[23-48.6]	39.7%[26.5-52.6]	24.6%[12.4-36.8]	23.3%[11.4-35.3]	20.8%[9.3-32.4]
	PTCy (n=161)	36%[28.5-43.6]	34.1%[25.9-42.4]	34%[25.6-42.5]	26.5%[19.6-34]	39.5%[30.7-48.3]	48.4%[39.4-57.3]	30.8%[22.5-39.1]
	<i>P</i> -value	0.82	0.77	0.52	<b>0.06</b>	0.008	0.001	0.03

Abbreviations: GVHD – graft-versus-host disease; NRM – non-relapse mortality; LFS – leukemia-free survival; OS – overall survival; GRFS – GVHD-free/relapse-free survival; ATG – anti-thymocyte globulin; PTCy – post-transplant cyclophosphamide

Supplemental Table 3. Subset analysis of post-transplant outcomes by ATG dose in available subjects (N=81), stratified by < or ≥ to median dose of 20 mg/Kg.

ATG dose (N=81)	100-day Acute GVHD II-IV	2-year Chronic GVHD	2-year Relapse	2-year NRM	2-year LFS	2-year OS	2-year GRFS
<20 mg/Kg	53.4% [33.5-69.8]	26.3% [9.8-46.4]	51.2% [25.3-72.1]	26% [10.9-44.1]	22.9% [2.7-43]	25.9% [5.7-46.2]	19.9% [0-40.3]
≥20 mg/Kg	23.3% [11.9-36.8]	22.5% [10.8-36.7]	44.6% [29-59]	35.3% [21.1-49.8]	20.2% [8-32.4]	24.8% [11.7-37.9]	18.1% [6.4-29.8]
<i>P</i> -value	<b>0.007</b>	0.68	0.95	0.67	0.54	0.78	0.40

Abbreviations: GVHD – graft-versus-host disease; NRM – non-relapse mortality; LFS – leukemia-free survival; OS – overall survival; GRFS – GVHD-free/relapse-free survival; ATG – anti-thymocyte globulin