

Monoclonal gammopathy of clinical significance: *in vivo* demonstration of the anti-thrombotic effect of an acquired anti-thrombin antibody

Berenice Schell,¹ Celine Desconclois,¹ Xavier Mariette,^{2,3} Cecile Goujard,^{3,4} Peter J. Lenting,^{3,5} Cecile V. Denis^{3,5} and Valerie Proulle^{1,3,5}

¹Service Hematologie Biologique, Hopital Bicetre, AP-HP, Le Kremlin-Bicetre; ²Service Rhumatologie, Hopital Bicetre, AP-HP, INSERM UMR_S U1184, Le Kremlin-Bicetre; ³Universite Paris-Saclay, Paris; ⁴Service Medecine Interne, Hopital Bicetre, AP-HP, Le Kremlin-Bicetre and ⁵INSERM UMR_S 1176, Le Kremlin-Bicetre, France

Correspondence: VALERIE PROULLE - valerie.proulle@aphp.fr

doi:10.3324/haematol.2019.242370

Supplementary Figure 1

A

Patient's blood results

PARAMETERS	VALUE	NORMAL RANGE
Serum creatinine	1.0 mg/dL	(0.6-1.2 mg/dL)
Calcium	10.4 mg/dL	(8.5-10.5 mg/dL)
Total serum protein	69 g/L	(60-83 g/L)
Serum albumin	39.3 g/L	(36-50 g/L)
Alpha 1 fraction	3.1 g/L	(2.2-4.0 g/L)
Alpha 2 fraction	6.8 g/L	(5.0-8.7 g/L)
Beta 1 fraction	3.5 g/L	(3.4-5.5 g/L)
Beta 2 fraction	13.7 g/L	(2.1-4.7 g/L)
Gamma fraction	2.6 g/L	(8.0-16.0 g/L)
Hemoglobin	14.9 g/dL	(13-17 g/dL)
Platelets	158 G/L	(150-400 G/L)
Leucocytes	3.27 G/L	(4-10 G/L)
Serum IgA	0.38 g/L	(1.02-3.25 g/L)
Serum IgG	14.4 g/L	(6.49-12.05 g/L)
Serum IgM	0.18 g/L	(0.51-1.45 g/L)
Serum free light kappa chains	6.3 mg/L	(3.3-19.4 mg/L)
Serum free light lambda chains	7.11 mg/L	(5.71-26.3 mg/L)
kappa/lambda ratio in serum	0.89	(0.26-1.65)
Urine free light kappa chains	6.68 mg/L	(1.35-24.19 mg/L)
Urine free light lambda chains	0.95 mg/L	(0.24-6.66 mg/L)
kappa/lambda ratio in urine	7.03	(2.04-10.37)
β2 microglobulin (serum)	2.13 mg/L	(1.09-2.53 mg/L)
Rheumatoid factor	<15 U/mL	(<15 U/mL)
Antinuclear antibodies	negative	

B

Patient's electrophoresis of seric proteins (Capillary, Sebia)

