

Pre-treatment maximum standardized uptake value predicts outcome after frontline therapy in patients with advanced stage follicular lymphoma

Paolo Strati,¹ Mohamed Amin Ahmed,¹ Nathan H. Fowler,¹ Loretta J. Nastoupil,¹ Felipe Samaniego,¹ Luis E. Fayad,¹ Fredrick B. Hagemester,¹ Jorge E. Romaguera,¹ Alma Rodriguez,¹ Michael Wang,¹ Jason R. Westin,¹ Chan Cheah,¹ Mansoor Noorani,¹ Lei Feng,² Richard E. Davis¹ and Sattva S. Neelapu¹

¹Department of Lymphoma and Myeloma, The University of Texas MD Anderson Cancer Center and ²Department of Biostatistics, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

©2020 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2019.230649

Received: June 28, 2019.

Accepted: October 4, 2019.

Pre-published: October 10, 2019.

Correspondence: SATTVA S. NEELAPU - sneelapu@mdanderson.org

Supplementary Data

Pre-Treatment Maximum Standardized Uptake Value Predicts Outcome after Frontline Therapy in Patients with Advanced Stage Follicular Lymphoma

Strati et al.

Supplementary Table. Analysis of multiple single unit increments of SUV_{max} and PFS in 346 patients included in the study. A starting threshold of 13 was selected, based on historical association between SUV_{max} and risk of transformation. No significant association between SUV_{max} thresholds and PFS could be observed when limiting the analysis to the 294 patients with SUV_{max} < 18

SUV _{max}	Patients (N=346)	Events	Median PFS (months)	HR [95% CI]	p-value
<= 13	224	85	Not reached	1 [0.73-1.5]	0.82
> 13	122	44	Not reached		
<= 14	240	90	145	1 [0.71-1.5]	0.85
> 14	106	39	Not reached		
<= 15	250	94	145	1 [0.69-1.5]	0.91
> 15	96	35	Not reached		
<= 16	270	100	Not reached	1.1 [0.74-1.7]	0.61
> 16	76	29	Not reached		
<= 17	283	104	Not reached	1.2 [0.80-1.9]	0.34
> 17	63	25	114		
<= 18	294	106	Not reached	1.5 [0.95-2.3]	0.08
> 18	52	23	114		
<= 19	302	111	Not reached	1.3 [0.79-2.2]	0.29
> 19	44	18	114		
<= 20	305	112	Not reached	1.4 [0.82-2.3]	0.23
> 20	41	17	114		