## **SUPPLEMENTARY APPENDIX**

# Entospletinib and obinutuzumab in patients with relapsed/refractory chronic lymphocytic leukemia and B-cell malignancies

Adam S. Kittai, <sup>1</sup> Scott Best, <sup>1</sup> Bria Thurlow, <sup>1</sup> Vi Lam, <sup>1</sup> Taylor Hashiguchi, <sup>1</sup> Shaun Goodyear, <sup>1</sup> Daniel O. Persky, <sup>2</sup> Craig Okada, <sup>1</sup> Byung Park, <sup>1</sup> Stephen E. Spurgeon <sup>1</sup> and Alexey V. Danilov <sup>1,3</sup>

<sup>1</sup>Knight Cancer Institute, Oregon Health & Science University, Portland, OR; <sup>2</sup>University of Arizona Cancer Center, Tucson, AZ and <sup>3</sup>City of Hope Comprehensive Cancer Center, Duarte, CA, USA

°Current address: Ohio State University, Columbus, OH, USA Correspondence: ALEXEY V. DANILOV - adanilov@coh.org

doi:10.3324/haematol.2020.270298

#### **SUPPLEMENTAL TABLE S1: PHASE 1 DOSING SCHEMA**

Dose Level (DL)	Entospletinib dose (starting on day -7 of run-in phase)	Obinutuzumab (Dosed cycles 1-6)		
-1	200 mg PO daily			
1	200 mg PO twice daily	C1D1: 100 mg IV C1 days 2, 8, 15: 1000 mg IV C2-C6: 1000 mg IV on day 1		
2	400 mg PO twice daily	g ,		

# SUPPLEMENTAL TABLE S2: CHARACTERISTICS OF PATIENTS ENROLLED ON STUDY

Characteristics	All patients			
Characteristics	N = 23	(%)		
Age				
< 65 years	8	(34.8)		
≥ 65 years	15	(65.2)		
Sex				
Male	17	(73.9)		
Female	6	(26.1)		
Race				
White	20	(87.0)		
Not Reported	3	(13.0)		
ECOG performance status				
0	9	(39.1)		
1	13	(56.5)		
2	1	(4.4)		
Dose Level 1	6	(26.1)		
Dose Level 2	17	(73.9)		
Chronic lymphocytic leukemia	21	(91.3)		
Follicular Lymphoma	2	(8.7)		

## SUPPLEMENTAL TABLE S3. EFFICACY IN PATIENTS WITH CLL

IWCLL Response after:	3 Cycles		6 Cycles		≥9 Cycles		Best Response	
	No.	%	No.	%	No.	%	No.	%
CR	0	0	2	9.5	3	14.3	3	14.3
PR	11	52.4	11	52.4	11	52.4	11	52.4
ORR	11	52.4	13	52.4	14	66.7	14	66.7