A phase I study of the fully human, fragment crystallizable-engineered, anti-CD-33 monoclonal antibody BI 836858 in patients with previously-treated acute myeloid leukemia

Sumithira Vasu,¹ Jessica K. Altman,² Geoffrey L. Uy,³ Martin S. Tallman,⁴ Ivana Gojo,⁵ Gerard Lozanski,⁶ Ute Burkard,⁷ Annika Osswald,⁷ Pamela James,⁸ Björn Rüter⁸ and William Blum⁹

¹Division of Hematology, Department of Internal Medicine, The Ohio State University Comprehensive Cancer Center, Columbus, OH, USA; ²Division of Hematology/Oncology, Northwestern University, Chicago, IL, USA; ³Washington University School of Medicine, Siteman Cancer Center, St Louis, MO, USA; ⁴Leukemia Service, Memorial Sloan Kettering Cancer Center, Weill Cornell Medical College, New York, NY, USA; ⁵The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, Baltimore, MD, USA; ⁶Department of Pathology, The Ohio State University, Columbus, OH, USA; ⁷Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach an der Riβ, Germany; ⁸Boehringer Ingelheim Pharmaceuticals, Inc, Ridgefield, CT, USA and ⁹Department of Hematology and Medical Oncology, Emory University School of Medicine, Winship Cancer Institute, Atlanta, GA, USA

Correspondence:

WILLIAM BLUM - william.g.blum@emory.edu doi:10.3324/haematol.2020.274118

Supplementary figures



Supplementary Figure 1: Comparison of individual and gMean of C_{max} (dose norm.) after single iv administration of 10 to 40 mg BI 836858 in Cycle 1 (1st and 2nd infusion)

Supplementary Figure 2: Correlation between Cmax of BI 836858 and CD33+ Blasts_ABS (in peripheral blood) after single iv administration of 10 to 40 mg BI 836858 in cycle 1 (1st and 2nd infusion)



Supplementary Figure 3. Proportion of NK cells over time in (A) peripheral blood and (B) bone marrow of patients with R/R AML treated with 40 mg BI 836858. R/R AML: relapsed or refractory acute myeloid leukemia; NK = natural killer; scr: screening; WBC: white blood cell



в

