

Dual pyroptotic biomarkers predict erythroid response in lower-risk non-del(5q) myelodysplastic syndromes treated with lenalidomide and recombinant erythropoietin

Chen Wang,^{1,2} Kathy L. McGraw,² Amy F. McLemore,² Rami Komrokji,² Ashley A. Basiorka,² Najla Al Ali,² Jeffrey E. Lancet,² Eric Padron,² Olivier Kosmider,³ Michaela Fontenay,³ Pierre Fenaux,⁴ Alan F. List^{2#} and David A. Sallman^{2#}

¹Department of Internal Medicine, University of South Florida, Morsani College of Medicine, Tampa, FL, USA; ²Department of Malignant Hematology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA; ³Service d'Hématologie Biologique, Assistance Publique-Hôpitaux de Paris. Centre-Université de Paris, Hôpital Cochin, Paris, France and ⁴Groupe Francophone des Myélodysplasies, Hôpital Saint Louis, Paris, France

#AFL and DAS contributed equally as co-senior authors.

Correspondence:

DAVID A. SALLMAN - david.sallman@moffitt.org

doi:10.3324/haematol.2021.278855

Supplement table. Patient's characteristics based on treatment

Median (range) or N (%)	All (n = 69)	Lenalidomide (n = 31)	Lenalidomide plus epoetin β (n = 38)	P*
<i>Demographics</i>				
Age (years)	73 (46-86)	73 (54-79)	74 (46-86)	0.089
Male (%)	48 (69.6)	19 (61.3)	29 (76.3)	0.177
<i>Clinical features</i>				
Hemoglobin (g/dL)	8.1 (6.3-10.0)	7.8 (6.8-9.6)	8.1 (6.3-10.0)	0.064
ANC (/mm ³)	2.1 (0.15-10.06) (n = 67)	2.00 (0.15-10.06) (n = 30)	2.60 (0.40-5.70) (n = 37)	0.695
Platelet (10 ³ /mm ³)	235 (58-587) (n = 67)	209 (68-494) (n = 30)	242 (58-587) (n = 37)	0.854
Serum EPO > 200 mU/mL (%)	27 (48.2) (n = 56)	12 (n = 29)	15 (n = 27)	0.289
Bone marrow blast > 2% (%)	27 (39.1)	8 (25.8)	19 (50.0)	0.041
Bone marrow erythroblast (%)	28.9 (1.0-79.0) (n = 67)	36.7 (1.0-65.0) (n = 30)	27.0 (4.4-79.0) (n = 37)	0.308
RARS (%)	38 (55.1)	17 (54.8)	21 (55.3)	0.972
Favorable karyotype (%)	59 (86.8) (n = 68)	27 (87.1)	32 (86.5) (n = 37)	1.000
<i>Splicesome mutation</i>				
<i>SF3B1</i> (%)	52 (75.4)	26 (83.9)	26 (68.4)	0.168
<i>U2AF</i> (%)	0 (0.0)	0 (0.0)	0 (0.0)	1.000
<i>SRSF2</i> (%)	6 (8.7)	4 (12.9)	2 (5.3)	0.397
<i>ZRSR2</i> (%)	5 (7.2)	2 (6.5)	3 (7.9)	1.000
<i>Epigenetic mutation</i>				
<i>TET2</i> (%)	34 (49.3)	14 (45.2)	20 (52.6)	0.537
<i>IDH</i> (%)	2 (2.9)	2 (6.5)	0 (0.0)	0.198
<i>DNMT3A</i> (%)	13 (18.9)	6 (19.4)	7 (18.4)	0.921
<i>ASXL1</i> (%)	12 (17.4)	8 (25.8)	4 (10.5)	0.119
<i>EZH2</i> (%)	2 (2.9)	0 (0.0)	2 (5.3)	0.498
Number of mutations	2 (0-6)	2 (1-6)	2 (0-6)	0.515
IPSS low risk (%)	33 (48.5) (n = 68)	16 (51.6)	17 (45.9) (n = 37)	0.641
<i>Biomarker</i>				
ASC specks > median (%)	34 (49.3)	11 (35.5)	23 (60.5)	0.039
<i>Treatment</i>				
Prior responder to recombinant EPO (%)	24 (35.8) (n = 67)	9 (29.0) (n = 31)	15 (41.7) (n = 36)	0.282
Responder (%)	28 (40.6)	7 (22.6)	21 (55.3)	0.006

* Continuous variables were compared by Mann-Whitney test, and categorical variables were compared using Fisher's exact test or chi-square test, as appropriate.

Abbreviations: ANC, absolute neutrophil count; EPO, erythropoietin; RARS, refractory anemia with ring sideroblasts; IPSS, international prognostic score system; ASC, apoptosis-associated speck-like protein containing a CARD.