

Full *versus* prophylactic-intermediate doses of anticoagulants in COVID-19: a meta-analysis

Lorenzo Loffredo,¹ Augusto Di Castelnuovo,² Giovanni Alfonso Chiariello,^{3,4} Pasquale Pignatelli¹ and Francesco Violi^{2,5}

¹Department of Clinical Internal Medicine, Anesthesiologic and Cardiovascular Sciences, Sapienza University of Rome, Rome;

²Mediterranea Cardiocentro, Napoli; ³Cardiovascular Sciences Department, Agostino Gemelli Foundation Polyclinic IRCCS, Rome;

⁴Catholic University of The Sacred Heart, Rome and ⁵Sapienza University of Rome, Rome, Italy.

Correspondence:

F. VIOLI - francesco.violi@uniroma1.it

L. LOFFREDO - lorenzo.loffredo@uniroma1.it

<https://doi.org/10.3324/haematol.2022.280652>

SUPPLEMENTARY DATA

Full versus prophylactic-intermediate doses of anticoagulants in COVID-19: a meta-analysis.

Lorenzo Loffredo¹, Augusto Di Castelnuovo², Giovanni Aldo Chiariello^{3,4}, Pasquale Pignatelli¹ and Francesco Violi^{1,2}.

¹Department of Clinical, Internal Medicine, Anesthesiologic and Cardiovascular Sciences Sapienza University of Rome, Rome, Italy.


²Mediterranea Cardiocentro, Napoli, Italy.

³Cardiovascular Sciences Department, Agostino Gemelli Foundation Polyclinic IRCCS, Rome, Italy

⁴Catholic University of The Sacred Heart, Rome, Italy.







Supplementary Figure 1

Flow-chart diagram



Supplementary Figure 2: Funnel Plots

Funnel Plot of Standard Error by Log risk ratio



Supplementary Figure 3: NNT and NNH.

