

Axicabtagene ciloleucel compared to tisagenlecleucel for the treatment of aggressive B-cell lymphoma

Mi Kwon,^{1,2*} Gloria Iacoboni,^{3,4*} Juan Luis Reguera,⁵ Lucía López Corral,⁶ Rafael Hernani Morales,⁷ Valentín Ortiz-Maldonado,⁸ Manuel Guerreiro,⁹ Ana Carolina Caballero,¹⁰ María Luisa Guerra Domínguez,¹¹ Jose María Sanchez Pina,¹² Alberto Mussetti,¹³ Juan Manuel Sancho,¹⁴ Mariana Bastos-Oreiro,^{1,2} Eva Catala,^{3,4} Javier Delgado,⁵ Hugo Luzardo Henriquez,¹¹ Jaime Sanz,⁹ María Calbacho,¹² Rebeca Bailén,^{1,2} Cecilia Carpio,^{3,4} Jose María Ribera,¹⁴ Anna Sureda,¹³ Javier Briones,¹⁰ Juan Carlos Hernandez-Boluda,⁷ Nuria Martínez Cebrián,⁸ Jose Luis Diez Martín,^{1,2,15} Alejandro Martín⁶ and Pere Barba^{3,4} on behalf of GETH-TC (Spanish Group of Stem Cell Transplantation and Cell Therapy) and GELTAMO (Spanish Group of Lymphoma and Autologous Stem Cell Transplantation)

Correspondence: M. Kwon
mi.kwon@salud.madrid.org

Received: February 7, 2022.

Accepted: June 17, 2022.

Prepublished: June 30, 2022.

<https://doi.org/10.3324/haematol.2022.280805>

©2023 Ferrata Storti Foundation

Published under a CC BY-NC license 

¹Department of Hematology, Hospital General Universitario Gregorio Marañón, Madrid;
²Instituto de Investigación Sanitaria Gregorio Marañón, Madrid; ³Department of Hematology, Vall d'Hebron University Hospital, Experimental Hematology, Vall d'Hebron Institute of Oncology (VHIO), Vall d'Hebron Barcelona Hospital Campus, Barcelona; ⁴Department of Medicine, Universitat Autònoma de Barcelona, Bellaterra; ⁵Department of Hematology, Hospital Universitario Virgen del Rocío, Instituto de Biomedicina de Sevilla, Sevilla;
⁶Department of Hematology, Hospital Clínico Universitario de Salamanca, IBSAL, Salamanca;
⁷Department of Hematology, Hospital Clínico Universitario de Valencia, Instituto de Investigación Sanitaria INCLIVA, Valencia; ⁸Department of Hematology, Hospital Clínic, Barcelona; ⁹Department of Hematology, Hospital Universitario y Politécnico La Fe, Valencia;
¹⁰Department of Hematology, Hospital de la Santa Creu i Sant Pau, Barcelona; ¹¹Department of Hematology, Hospital Universitario de Gran Canaria Doctor Negrín, Las Palmas de Gran Canaria; ¹²Department of Hematology, Hospital Universitario 12 de Octubre, Madrid;
¹³Department of Hematology, Hospital Duran i Reynals, Instituto Catalán de Oncología, Barcelona; ¹⁴Department of Hematology, Hospital Universitari Germans Trias i Pujol, Instituto Catalán de Oncología, Josep Carreras Research Institute, Badalona and ¹⁵Universidad Complutense de Madrid, Madrid, Spain

*MK and GI contributed equally as co-first authors.

Supplemental Material

Table S1. Characteristics of infused patients before lymphodepletion

	Total infused n=261	Axi-cel infused n=134	Tisa-cel infused n=127	p
Bridging therapy, n (%)	210 (80) 1 Line 2 Lines	104 (78) 95 (71) 9 (7)	106 (83) 99 (78) 7 (5)	0.275
ECOG, n (%)	244 (94) 0-1 2-4	123 (92) 11 (8)	121 (95) 6 (5)	0.319
R-IPI prognostic score, n (%)	87 (33) 0-2 95 (36) 3-5 79 (30) NA	51 (38) 57 (43) 26 (19)	36 (28) 38 (30) 53 (42)	0.881
Bulky disease, n (%)	67 (26)	41 (31)	26 (20)	0.248
Extranodal disease, n (%)				
≥ 2 sites	47 (18)	30 (22)	17 (14)	0.051
Disease status, n (%)				
Progressive disease	149 (57)	81 (60)	68 (54)	
Stable disease	43 (16)	24 (18)	19 (15)	0.625
Partial response	25 (10)	15 (11)	10 (8)	
Complete response	11 (4)	4 (3)	7 (5)	
NA	33 (13)	10 (8)	23 (18)	
LDH >ULN, n (%)	143 (55)	67 (50)	76 (60)	0.440
CRP >ULN, n (%)	154 (59)	71 (53)	83 (65)	0.119
Platelets $\times 10^3/\mu\text{L}$				
Median (range)	141.5 (119-558)	142 (28-550)	141 (19-558)	0.921

Abbreviations: R-IPI, revised international prognostic index; NA, not available; LDH, lactate dehydrogenase, CRP, c-reactive protein; >ULN, upper limit of normal

Tables S2.

A. Bridging therapy regimens used in axicabtagene ciloleucel and tisagenlecleucel–treated patients.

	Total infused n=261	Axi-cel infused n=134	Tisa-cel infused n=127
Bridging therapy, n (%)	210 (80)	104 (78)	106 (83)
Chemotherapy*	127 (48)	64 (48)	63 (49)
Low intensity regimen&	43 (16)	12 (9)	31 (24)
Radiotherapy	18 (7)	13 (10)	5 (4)
Steroids only	12 (5)	10 (7)	2 (1.5)
Other\$	4 (1.5)	1 (0.7)	3 (2)
Unknown	6 (2)	4 (3)	2 (2)

*: Gemcitabine based n=67, intensive regimen n=27 (R-ICE n=13, R-EISHAP or DHAP n= 3, R-MINE n=3, reduced R-HyCVAD n=1, reduced BEAM n=1, Burkimab n=1, R-MACOB n=1, other n=4), bendamustine based n=22 (11 combined with polatuzumab), cyclophosphamide based n=11

&: Cyclophosphamide +/- prednisone +/- rituximab n=34, R-lenalidomide n=5, rituximab n=1, polatuzumab + rituximab n=1, ibrutinib n=1, brentuximab n=1

\$: pembrolizumab n=1, pixantrona n=3

B. Response rates reported to the different bridging therapies.

	N	Complete Response	Partial Response	Stable Disease	Progressive Disease
Chemotherapy	116	7 (6)	15 (12)	16 (14)	68 (59)
Low intensity regimen	42	1 (2)	3 (7)	4 (10)	22 (52)
Polatuzumab-based					
PBR	11	2 (18)	3 (27)	2 (18)	4 (37)
PR	1	0 (0)	0 (0)	0 (0)	1 (100)
Radiotherapy	18	0 (0)	1 (6)	4 (22)	13 (72)
Steroids	12	0 (0)	0 (0)	1 (8)	11 (92)
Other/unkwnown	10	0 (0)	0 (0)	0 (0)	7 (70)

PBR: polatuzumab-bendamustine-rituximab; PR: polatuzumab-rituximab

Table S3. A. Univariable analysis for PFS and OS. B. Logistic regression analysis for CRS grade ≥3 and ICANS grade ≥3.

A. Univariable analysis for PFS and OS

	PFS		OS	
	HR (95%CI)	p	HR (95%CI)	p
<i>CART type. axi-cel vs tisa-cel</i>	0.781 (0.586-1.040)	0.090	0.735 (0.527-1.024)	0.069
<i>Sex. male vs female</i>	0.992 (0.737-1.335)	0.958	1.025 (0.731-1.437)	0.885
<i>Age</i>	0.994 (0.982-1.007)	0.385	1.000 (0.984-1.015)	0.946
<i>HCT-CI. 0-2 vs >2</i>	0.930 (0.653-1.323)	0.685	0.944 (0.644-1.385)	0.770
<i>Histology DLBCL vs other</i>	1.222 (0.881-1.694)	0.229	1.067 (0.745-1.528)	0.723
<i>Cell of origin. GCB vs non-GCB</i>	0.780 (0.572-1.063)	0.116	0.754 (0.532-1.067)	0.111
<i>Number prior lines. 2 vs >2</i>	1.088 (0.946-1.251)	0.235	1.134 (0.974-1.321)	0.106
<i>Primary Refractory. yes vs no</i>	1.440 (1.065-1.946)	0.018	1.647 (1.161-2.336)	0.005
<i>Previous ASCT. yes vs no</i>	0.608 (0.438-0.843)	0.003	0.470 (0.310-0.713)	0.000
<i>Disease status. PD vs other</i>	2.179 (1.361-3.490)	0.001	3.803 (1.728-8.372)	0.001
<i>ECOG at apheresis 0-1 vs >1</i>	2.529 (1.336-4.786)	0.004	3.885 (2.034-7.419)	0.000
<i>Disease stage I-II vs III-IV</i>	1.773 (1.26-2.496)	0.001	1.896 (1.226-2.841)	0.002
<i>R-IPI at apheresis</i>	1.291 (1.134-1.470)	0.000	1.535 (1.316-1.791)	0.000
<i>Bulky size at apheresis. yes vs no</i>	1.810 (1.322-2.479)	0.000	1.989 (1.397-2.831)	0.000
<i>LDH at apheresis</i>	2.444 (1.738-3.437)	0.000	3.140 (2.051-4.807)	0.000
<i>CRP at apheresis</i>	2.212 (1.54-3.178)	0.000	2.478 (1.644-3.735)	0.000
<i>Platelets at apheresis</i>	0.998 (0.996-1.000)	0.061	0.996 (0.993-1.000)	0.026
<i>Bridging therapy. yes vs no</i>	1.594 (1.023-2.484)	0.040	2.665 (1.423-4.993)	0.002
<i>Disease status at LD. PD vs other</i>	2.797 (1.891-4.139)	0.000	2.500 (1.508-4.142)	0.000
<i>Bulky size at LD. yes vs no</i>	1.946 (1.386-2.733)	0.000	1.916 (1.264-2.903)	0.002
<i>Extranodal at LD >2 sites. yes vs no</i>	1.450 (1.297-1.622)	0.000	1.441 (1.262-1.646)	0.000
<i>ECOG at LD. 0-1 vs >1</i>	3.056 (1.572-5.944)	0.001	3.072 (1.600-5.897)	0.001
<i>R-IPI at LD</i>	1.301 (1.152-1.470)	0.000	1.343 (1.142-1.580)	0.000
<i>Platelets at LD</i>	0.999 (0.997-1.001)	0.269	0.997 (0.995-1.000)	0.063
<i>LDH at LD</i>	2.233 (1.586-3.144)	0.000	2.598 (1.669-4.043)	0.000
<i>CRP at LD</i>	2.495 (1.644-3.786)	0.000	3.190 (1.877-5.422)	0.000
<i>LDH pre infusion</i>	2.179 (1.463-3.245)	0.000	2.175 (1.324-3.571)	0.002
<i>CRP pre infusion</i>	2.664 (1.570-4.521)	0.000	4.497 (2.128-9.5)	0.000
<i>CRS. yes vs no</i>	0.740 (0.514-1.064)	0.104	1.071 (0.665-1.727)	0.777
<i>CRS 3-4. yes vs no</i>	1.533 (0.752-3.126)	0.239	2.615 (1.250-5.470)	0.011
<i>CRS tocilizumab. yes vs no</i>	0.886 (0.644-1.218)	0.455	1.337 (0.907-1.971)	0.143
<i>CRS steroids. yes vs no</i>	0.895 (0.584-1.372)	0.611	1.219 (0.721-2.062)	0.460
<i>ICANS. yes vs no</i>	0.974 (0.679-1.396)	0.884	1.450 (0.942-2.234)	0.092
<i>ICANS 3-4. yes vs no</i>	1.108 (0.620-1.978)	0.730	1.690 (0.842-3.392)	0.140
<i>ICANS tocilizumab. yes vs no</i>	0.896 (0.579-1.388)	0.623	1.475 (0.884-2.462)	0.137
<i>ICANS steroids. yes vs no</i>	0.990 (0.660-1.483)	0.960	1.396 (0.846-2.301)	0.191

B. Univariable logistic regression analysis for CRS grade ≥3 and ICANS grade ≥3

	CRS grade ≥ 3		ICANS grade ≥ 3	
	OR (95%CI)	P	OR (95%CI)	P
<i>CART type. axi-cel vs tisa-cel</i>	1.330 (0.516-3.429)	0.555	4.400 (1.731-11.180)	0.002
<i>Age</i>	1.001 (0.995-1.050)	0.970	1.004 (0.968-1.041)	0.836
<i>HCT-CI. 0-2 vs >2</i>	0.401 (0.090-1.795)	0.232	1.106 (0.447-2.733)	0.828
<i>Histology DLBCL vs other</i>	0.719 (0.395-1.309)	0.281	0.914 (0.609-1.372)	0.665
<i>Cell of origin. GCB vs non-GCB</i>	0.671 (0.239-1.882)	0.448	1.438 (0.572-3.615)	0.440
<i>Number prior lines. 2 vs ></i>	1.067 (0.676-1.684)	0.782	0.534 (0.336-0.848)	0.008
<i>Primary Refractory. yes vs o</i>	1.270 (0.482-3.345)	0.628	1.105 (0.508-2.404)	0.801
<i>LDH at apheresis</i>	2.547 (0.817-7.938)	0.107	0.984 (0.438-2.206)	0.968
<i>CRP at apheresis</i>	1.779 (0.537-5.895)	0.346	0.539 (0.228-1.271)	0.158
<i>Leucocyte count at apheresis</i>	0.898 (0.701-1.150)	0.392	1.040 (0.940-1.151)	0.449
<i>Lymphocyte count at apheresis</i>	0.922 (0.581-1.463)	0.730	1.020 (0.774-1.346)	0.886
<i>Platelet count at apheresis</i>	0.993 (0.984-1.002)	0.121	1.000 (0.997-1.003)	0.927
<i>Bridging therapy. yes vs no</i>	2.158 (0.481-9.683)	0.315	2.361 (0.685-8.131)	0.173
<i>Disease status PD vs other</i>	1.296 (0.439-3.830)	0.639	1.219 (0.504-2.950)	0.660
<i>Bulky at LD. yes vs no</i>	1.985 (0.760-5.185)	0.162	1.250 (0.534-2.925)	0.608
<i>Extranodal at LD. 0-2 vs >2</i>	1.012 (0.7701-331)	0.930	1.225 (0.890-1.686)	0.213
<i>ECOG at LD. 0-1 vs >1</i>	6.756 (2.082-21.920)	0.001	2.658 (0.803-8.800)	0.109
<i>R-IPI at LD. 0-2 vs >2</i>	2.717 (0.829-8.907)	0.099	2.315 (0.901-5.947)	0.081
<i>Platelets at LD</i>	1.001 (0.996-1.006)	0.682	1.002 (0.992-1.006)	0.337
<i>LDH at LD</i>	6.299 (1.411-28.120)	0.016	0.924 (0.426-2.001)	0.840
<i>CRP at LD</i>	2.705 (0.761-9.624)	0.124	1.961 (0.760-5.057)	0.164
<i>LDH at infusion</i>	1.026 (0.236-4.459)	0.973	1.461 (0.569-3.749)	0.430
<i>CRP at infusion</i>	2.162 (0.480-9.740)	0.315	1.591 (0.524-4.832)	0.413

Abbreviations: HCT-CI, Hematopoietic Cell Transplantation-Comorbidity Index; GCB, germinal center B-cell; ASCT, autologous stem cell transplantation; PD, progressive disease; R-IPI, revised international prognostic index; LDH, lactate dehydrogenase; CRP, c-reactive protein; CRS, cytokine release syndrome; ICANS, immune effector cell-associated neurotoxicity syndrome; LD, lymphodepletion.

Table S4. Infections within the first six months after infusion in axicabtagene ciloleucel and tisagenlecleucel–treated patients.

	Total infused (n=261)	Axi-cel infused (n=134)	Tisa-cel infused (n=127)
Infectious episodes, n	91	56	35
Bacterial, n (%)	54 (59)	32 (57)	22 (63)
Mild	3 (3)	1 (2)	2 (6)
Moderate	9 (11)	3 (5)	6 (17)
Severe	37 (41)	25 (45)	12 (34)
Life-threatening	4 (4)	2 (4)	2 (6)
Fatal	1 (1)	1 (2)	0 (0)
Viral, n (%)	31 (34)	22 (36)	9 (26)
Mild	13 (14)	7 (13)	6 (17)
Moderate	6 (7)	4 (7)	2 (6)
Severe	5 (5)	4 (7)	1 (3)
Life-threatening	6 (7)	6 (11)	0 (0)
Fatal	1 (1)	1 (2)	0 (0)
Fungal, n (%)	6 (7)	2 (4)	4 (11)
Mild	1 (1)	0 (0)	1 (3)
Moderate	1 (1)	1 (2)	0 (0)
Severe	2 (2)	0 (0)	2 (6)
Life-threatening	1 (1)	0 (0)	1 (3)
Fatal	1 (1)	1 (2)	0 (0)

Figure S1. Reasons for not undergoing CAR T-cell infusion for patients intended to be treated with axi-cel and tisa-cel





Figure S2. Timeline of neurotoxicity (ICANS) after infusion of tisagenlecleucel (A) and axicabtagene ciloleucel (B).

A



B




Figure S3. Non-relapse mortality causes in axicabtagene ciloleucel and tisagenlecleucel–treated patients






Figure S4. Response assessment in axicabtagene ciloleucel and tisagenlecleucel–treated patients at days 30, 100 and 180 after infusion.




Abbreviations: CR, complete response; PR, partial response; SD: stable disease; PD: progressive disease

Figure S5. Progression free survival (PFS) and Overall survival (OS) from apheresis of patients intended to be treated with axicabtagene ciloleucel and tisagenlecleucel.

A



B



Number at risk

Tisagenlecleucel	155	142	130	98	85	58	48	41	38	35	30	26	25	24	23	21
Axi-cel	152	147	133	108	90	65	54	53	44	33	30	26	25	20	16	12

155	150	137	121	109	91	83	71	66	62	56	50	46	41	36	30
152	148	135	108	113	94	81	75	67	58	48	41	34	29	24	19