

An open-label, phase I/II trial to determine the maximum tolerated dose and investigate safety, pharmacokinetics and efficacy of BI 836858, an unconjugated anti-CD33 monoclonal antibody, in combination with decitabine in patients with acute myeloid leukemia

Walter Fiedler,¹ Pau Montesinos,^{2,3} Christoph Schliemann,⁴ Jan Middeke,⁵ Sumithira Vasu,⁶ Christian W. Scholz,⁷ Jordi Esteve,⁸ Shoubhik Mondal,⁹ Björn Rüter,¹⁰ Ute Burkard,¹⁰ Annika Osswald¹⁰ and William Blum¹¹

¹Department of Oncology, Hematology and Bone Marrow Transplantation with Section Pneumology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; ²Hospital Universitari i Politècnic La Fe, Valencia, Spain; ³CIBERONC, Instituto Carlos III, Madrid, Spain; ⁴Department of Medicine A, Hematology and Oncology, University Hospital Muenster, Muenster, Germany; ⁵Uniklinikum Dresden, Dresden, Germany; ⁶Division of Hematology, The Ohio State University Comprehensive Cancer Center, Columbus, OH, USA; ⁷Department of Hematology and Oncology, Vivantes Klinikum Am Urban, Berlin, Germany; ⁸Hematology Department, Hospital Clínic of Barcelona, IDIBAPS, University of Barcelona, Barcelona, Spain; ⁹Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, CT, USA; ¹⁰Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach/Riss, Germany and ¹¹Department of Hematology and Medical Oncology, Winship Cancer Institute, Emory University School of Medicine, Atlanta, GA, USA.

Correspondence:


W. BLUM - william.g.blum@emory.edu

<https://doi.org/10.3324/haematol.2022.281128>

Supplementary Table S1. Comparison of Cmax in cycle 1, day 9 and day 23, for all dose groups.

Plasma concentrations		
Dose	Day 9, 6 h	Day 23, 6 h
[mg]	[ng/mL]	[ng/mL]
20	1790	4390
40	6400	7210
80	14,700	22,300

Supplementary Figure S1. Percentage CD33+ blasts (A), CD3-/CD16+ NK cells (B) and CD3-/CD16+/CD69+ activated NK cells (C) in the peripheral blood of individual patients treated with treated with BI 836858 80 mg.



NK: natural killer.