

# Response-adjusted regimen combining ruxolitinib, etoposide and dexamethasone (adRED) in adult patients with acute myeloid leukemia-associated hemophagocytic lymphohistiocytosis: a single-center pilot trial

Grégoire Stalder,<sup>1,2,3</sup> Madeleine Suffiotti,<sup>4</sup> Amandine Segot,<sup>1</sup> Alessandra Noto,<sup>4</sup> Giuseppe Pantaleo,<sup>3,4</sup> Olivier Spertini<sup>1,3</sup> and Michel Obeid<sup>3,4</sup>

<sup>1</sup>Service and Central Laboratory of Hematology, Lausanne University Hospital, Lausanne; <sup>2</sup>Service of Hematology and Laboratory of Hematology, Institut Central des Hôpitaux, Hôpital du Valais, Sion and <sup>3</sup>University of Lausanne, Lausanne and <sup>4</sup>Division of Immunology and Allergy, Lausanne University Hospital, Lausanne, Switzerland

Correspondence:  
M. OBEID - michel.obeid@chuv.ch

<https://doi.org/10.3324/haematol.2022.281221>

## Supplement document

<b>Supplemental Table 1.</b> Patient's and hemophagocytic lymphohistiocytosis diagnosis characteristics .....	2
<b>Supplemental Figure 1.</b> Differences in the classic parameter between time points of interest at day 0 and day 7 in patients treated with adRED and historical control patients. ....	5
<b>Supplemental Figure 2.</b> Intergroup treatment type data comparisons of low/high doses of etoposide/DEX for patients receiving ruxolitinib were performed via Fisher's exact test. Low-dose etoposide regimen = cumulative dose $\leq$ 350 mg/m <sup>2</sup> . High-dose etoposide = cumulative dose >350 mg/m <sup>2</sup> . Low-dose DEX regimen = cumulative dose $\leq$ 550 mg. High-dose DEX = cumulative dose >550 mg.....	6

**Supplemental Table 1.** Patient's and hemophagocytic lymphohistiocytosis diagnosis characteristics


	All patients (n=6)	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patient 6
Age (Years)	52 (mean)	61	40	34	72	42	64
Sex	4M /2F	Male	Male	Female	Female	Male	Male
ELN AML risk category							
Intermediate	1 (17%)	0	1	0	0	0	0
Adverse	5 (83%)	1	0	1	1	1	1
Karyotype							
Normal	2 (33%)	0	1	0	0	1	0
Complex	3 (50%)	1	0	0	1	0	1
NGS							
FTL3-ITD	2 (33%)	0	1	0	0	1	0
TP53 mutation	2 (33%)	1	0	0	1	0	0
DNMT3A	1 (17%)	0	1	0	0	0	0
TET2	1 (17%)	0	1	0	0	0	0
ASXL1	1 (17%)	0	0	0	1	0	0
Timing of HLH diagnosis							
At AML diagnosis	2 (33%)	1	0	0	0	0	1
At AML relapse	4 (67%)	0	1	1	1	1	0
After allo-HSCT	2 (33%)	0	1	1	0	0	0

Before intensive chemotherapy	2 (33%)	1	1	0	0	0	0
After intensive chemotherapy	4 (67%)	0	0	1	1	1	1
During C1 chemotherapy	5 (83%)	1	1	1	1	0	1
During C2 chemotherapy	1 (17%)	0	0	0	0	1	0
Type of chemotherapy							
		CLAG-IDA	CLAG	FLAG	7+3	7+3 followed by FLAG	FLAG-IDA
Clinical parameters							
Fever (max. temperature)	6 (100%) (mean 39.3°C)	Yes (39°C)	Yes (39.8°C)	Yes (40°C)	Yes (38.5°C)	Yes (38.9°C)	Yes (39.7°C)
Organomegaly	5/6 (83%)	1	1	1	0	1	1
Laboratory parameters							
Hemophagocytosis on bone marrow biopsy	3 (50%)	0	0	0	1	1	1
Ferritin (µg/L)	37689 (mean)	5962	9250	17394	134490	35728	23309
sCD25 (pg/mL)	7471 (mean)	1196	8182	4612	5551	22010	3273

Triglyceride (mmol/L)	2.2 (mean)	2	1.5	1.9	3.3	3.1	1.5
Fibrinogen (g/L)	5.8 (mean)	5.5	2.4	6.7	1.7	9.4	9.1
ASAT (U/L)	181 (mean)	235	14	42	286	202	309
Hemoglobin (g/L)	85 (mean)	74	71	84	88	107	86
Leukocytes (G/L)	2.2 (mean)	4.6	1.3	<0.1	0.7	<0.1	<0.1
Neutrophils (G/L)	1.57 (mean)	3.45	1.01	<0.1	0.24	<0.1	<0.1
Thrombocytes (G/L)	28 (mean)	67	7	26	28	12	25
HScore	247 (mean)	206	263	214	263	261	272
HLH-2004 criteria	-	5/8	5/8	5/8	5/8	6/8	6/8

Allo-HSCT = allogeneic hematopoietic stem cell transplantation. AML = acute myeloid leukemia.

**Supplemental Figure 1.** Differences in the classic parameter between time points of interest at day 0 and day 7 in patients treated with adRED and historical control patients.



**Supplemental Figure 2.** Correlations heatmap between cytokines and classic parameters of patients with acute leukemias and HLH at follow-up. Color scale (blue-to-red) represents Pearson's correlation coefficient.

