

Allogeneic hematopoietic cell transplant for hairy cell leukemia: EBMT experience

Dai Chihara,¹ Luuk Gras,² Nienke Zinger,³ Nicolaus Kröger,⁴ Jiri Mayer,⁵ Jakob Passweg,⁶ Régis Peffault de Latour,⁷ Jenny Byrne,⁸ William Krüger,⁹ Jan-Paul Bohn,¹⁰ Uwe Platzbecker,¹¹ Igor Wolfgang Blau,¹² Francesca Bonifazi,¹³ Grzegorz Helbig,¹⁴ Andrew McDonald,¹⁵ Martin Mistrik,¹⁶ Mohamad Mohty,¹⁷ Ron Ram,¹⁸ Jaime Sanz,¹⁹ Carlos Vallejo Llamas,²⁰ Robert J. Kreitman,²¹ Patrick J. Hayden,²² Donal McLornan,²³ Olivier Tournilhac,²⁴ Michel van Gelder²⁵ and Ibrahim Yakoub-Agha²⁶

¹Department of Lymphoma and Myeloma, The University of Texas MD Anderson Cancer Center, Houston, TX, USA; ²EBMT Statistical Unit, Leiden, the Netherlands; ³EBMT Study Unit, Leiden, the Netherlands; ⁴University Hospital Eppendorf, Hamburg, Germany; ⁵University Hospital Brno, Brno, Czech Republic; ⁶University Hospital Basel, Basel, Switzerland; ⁷Hopital St. Louis 207, Paris, France; ⁸Nottingham University, Nottingham, UK; ⁹Klinik fuer Innere Medizin C, Greifswald, Germany; ¹⁰University Hospital Innsbruck, Innsbruck, Austria; ¹¹Medical Clinic and Polyclinic 1, Leipzig, Germany; ¹²Medizinische Klinik m. S. Hämatologie, Onkologie und Tumorimmunologie, Berlin, Germany; ¹³IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy; ¹⁴Silesian Medical University, Katowice, Poland; ¹⁵Alberts Cellular Therapy, Netcare Pretoria East Hospital, Pretoria, South Africa; ¹⁶University Hospital, Bratislava, Slovak Republic; ¹⁷Sorbonne University, Saint-Antoine Hospital, AP-HP, and INSERM UMRs 938, Paris, France; ¹⁸Tel Aviv Sourasky Medical Center and Sourasky Faculty of Medicine, Tel Aviv, Israel; ¹⁹University Hospital La Fe, Valencia, Spain; ²⁰Hospital Universitario Donostia, S Sebastian, Spain; ²¹Laboratory of Molecular Biology, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA; ²²Department of Haematology, Trinity College Dublin, St. James's Hospital, Dublin, Ireland; ²³University College London Hospitals NHS Trust, London, UK; ²⁴Department of Hematology and Cell Therapy, CHU Estaing, Université Clermont Auvergne, Clermont-Ferrand, France; ²⁵University Hospital Maastricht, Maastricht, the Netherlands and ²⁶CHU de Lille, Univ Lille, INSERM U1286, Lille, France

Correspondence:

D. CHIHARA - DChihara@mdanderson.org

O. TOURNILHAC - otournilhac@chu-clermontferrand.fr

<https://doi.org/10.3324/haematol.2022.281754>

Supplemental table 1. Treatment list in each line. Data was available for 11 patients.

		N (%)
1st line	2-CdA/Cladribine	6 (54.5)
	Interferon-a	3 (27.3)
	Interferon	1 (9.1)
	CHOP	1 (9.1)
2nd line	2-CdA/Cladribine	4 (44.4)
	Interferon	1 (11.1)
	Pentostatin	1 (11.1)
	Rituximab (CD20)	1 (11.1)
	SC	1 (11.1)
	2-CdA/Cladribine + Amsacrine	1 (11.1)
3rd line	2-CdA/Cladribine	2 (22.2)
	Interferon-a	1 (11.1)
	Nipent	1 (11.1)
	Pentostatin	1 (11.1)
	Rituximab (CD20)	3 (33.3)
	VACOP-B	1 (11.1)
4th line	2-CdA/Cladribine	1 (11.1)
	Interferon-a	2 (22.2)
	Interferon	1 (11.1)
	Rituximab (CD20)	3 (33.3)
	Pentostatin + Rituximab	1 (11.1)
	DEXA-BEAM	1 (11.1)
5th line	2-CdA/Cladribine	2 (25.0)
	Interferon-a	1 (12.5)
	Interferon	1 (12.5)
	Prednisone	1 (12.5)
	Vemurafenib	1 (12.5)
	Bendamustin + Rutiximab	1 (12.5)
	R-CHOP	1 (12.5)
6th line	2-CdA/Cladribine	1 (14.3)
	Clexane	1 (14.3)
	Ibrutinib	1 (14.3)
	Rituximab (CD20)	1 (14.3)
	Vemurafenib	1 (14.3)
	R-DHAX	1 (14.3)
	VACOP-B	1 (14.3)
7th line	Ofatumumab	1 (50.0)
	Rituximab (CD20)	1 (50.0)
	Ofatumumab	1 (100.0)
9th line	Dabrafenib + Trametinib	1 (100.0)
10th line	Nivolumab	1 (100.0)