

Long-term analysis of the RiBVD phase II trial reveals the unfavorable impact of *TP53* mutations and hypoalbuminemia in older adults with mantle cell lymphoma; for the LYSA group

Sylvain Carras,^{1,2} Alexia Torroja,² Anouk Emadali,³ Emilie Montaut,³ Nicolas Daguindau,⁴ Adrian Tempescul,⁵ Anne Moreau,⁶ Emmanuelle Tchernonog,⁷ Anna Schmitt,⁸ Roch Houot,⁹ Caroline Dartigeas,¹⁰ Sarah Barbieux,¹¹ Selim Corm,¹² Anne Banos,¹³ Ludovic Fouillet,¹⁴ Jehan Dupuis,¹⁵ Margaret Macro,¹⁶ Joel Fleury,¹⁷ Fabrice Jardin,¹⁸ Clementine Sarkozy,¹⁹ Ghandi Damaj,²⁰ Pierre Feugier,²¹ Luc Matthieu Fornecker,²² Cecile Chabrot,²³ Veronique Dorvaux,²⁴ Krimo Bouabdallah,²⁵ Sandy Amorim,²⁶ Reda Garidi,²⁷ Laurent Voillat,²⁸ Bertrand Joly,²⁹ Nadine Morineau,³⁰ Marie Pierre Moles,³¹ Hacene Zerazhi,³² Jean Fontan,³³ Yazid Arkam,³⁴ Magda Alexis,³⁵ Vincent Delwail,³⁶ Jean Pierre Vilque,³⁷ Loic Ysebaert,³⁸ Barbara Burroni,³⁹ Mary Callanan,⁴⁰ Steven Le Gouill¹⁹ and Rémy Gressin;² for the Lymphoma Study Association

Correspondence: S. Carras
scarras@chu-grenoble.fr

Received: June 29, 2023.
Accepted: November 22, 2023.
Early view: November 30, 2023.


<https://doi.org/10.3324/haematol.2023.283724>

©2024 Ferrata Storti Foundation
Published under a CC BY-NC license 

¹Université Grenoble Alpes, University Hospital, Institute For Advanced Biosciences (INSERM U1209, CNRSéUMR 5309, UGA), Molecular Biology Department, Grenoble; ²Université Grenoble Alpes, University Hospital, Institute For Advanced Biosciences (INSERM U1209, CNRS UMR 5309, UGA), Oncohematology Department, Grenoble; ³Université Grenoble Alpes, University Hospital, Institute for Advanced Biosciences (INSERM U1209, CNRS UMR 5309, UGA), Research & Innovation Unit, Grenoble; ⁴Hematology Department, Annecy Hospital, Annecy; ⁵Hematology Unit, Brest University Hospital, Brest; ⁶Pathology Department, University Hospital, Nantes; ⁷Hematology Department, University Hospital, Montpellier; ⁸Hematology Department, Cancerology Institute Bergonie, Bordeaux; ⁹Hematology Department, University Hospital, Rennes; ¹⁰Hematology Department, University Hospital, Tours; ¹¹Hematology Department, Dunkerque Hospital, Dunkerque; ¹²Hematology Department, Chambery Hospital, Chambery; ¹³Hematology Department, Bayonne Cote Basque Hospital, Bayonne; ¹⁴Hematology Department, University Hospital, Saint Etienne; ¹⁵Lymphoid Malignancies Unit, Henri Mondor University Hospital, Assistance Publique-Hôpitaux de Paris, Créteil; ¹⁶IHBN – Hematology Department, University Hospital, Caen; ¹⁷Hematology Department, Cancerology Institute, Clermont-Ferrand; ¹⁸Hematology Department, Centre Henri Becquerel, Rouen; ¹⁹Institut Curie, Paris and Paris Saint Quentin University, UVSQ, Paris; ²⁰Hematology Department, University Hospital, Caen; ²¹Hematology Department, University Hospital, Nancy; ²²Hematology Department, University Hospital, Strasbourg; ²³Hematology Department, University Hospital, Clermont-Ferrand; ²⁴Hematology Department, University Hospital, Metz; ²⁵Hematology Department, University Hospital, Bordeaux; ²⁶Hematology & Cellular Therapy Department, Hospital Saint Vincent de Paul, Université Catholique de Lille, Lille; ²⁷Hematology Department, Hospital Saint Quentin, Saint Quentin; ²⁸Hematology Department, Hospital Chalon sur Saone, Chalon sur Saone; ²⁹Hematology Department, Corbeil Hospital, Corbeil-Essonnes; ³⁰Hematology Department, Hospital La Roche Sur Yon, La Roche Sur Yon; ³¹Hematology Department, University Hospital Angers, Angers; ³²Hematology Department, Avignon Hospital, Avignon; ³³Hematology Department, University Hospital, Besançon; ³⁴Hematology Department, Mulhouse Hospital, Mulhouse; ³⁵Hematology Department, Orleans Hospital, Orleans; ³⁶Onc-Hematology Department, University Hospital Poitiers and INSERM, CIC 1402, University of Poitiers, Poitiers; ³⁷Hematology Department, Cancer Center Baclesse Caen, Caen; ³⁸Institut Universitaire du Cancer de Toulouse Oncopole, Toulouse; ³⁹Assistance Publique – Hôpitaux de Paris (APHP), Hôpital Cochin, Department of Pathology, Centre de Recherche des Cordeliers, Sorbonne University, Inserm, UMRS 1138, Université Paris Cité, F-75006 Paris and ⁴⁰Unit For Innovation in Genetics and Epigenetics and Oncology, Dijon University Hospital, Dijon, France

Supplementary Table 1: Demographic characteristics of the TP53mt and the TP53wt groups. 1 Median (IQR); n (%), 2 Wilcoxon rank sum test; Pearson's Chi-squared test

Characteristics	TP53 mt, N = 12 ¹	TP53 wt, N = 42 ¹	p-value ²
Age	73.0 (70.2, 76.0)	75.0 (69.0, 77.0)	0.8
Sex			>0.9
F	4 (33%)	16 (38%)	
M	8 (67%)	26 (62%)	
B symptoms			0.5
no	10 (91%)	31 (76%)	
yes	1 (9.1%)	10 (24%)	
Histology			>0.9
Blastoid/pleiomorphic	2 (18%)	7 (18%)	
Classic	9 (82%)	33 (82%)	
MIPI score			>0.9
high	10 (83%)	33 (79%)	
low/int	2 (17%)	9 (21%)	
KI67 (%)			0.027
<=30	0 (0%)	16 (47%)	
>30	9 (100%)	18 (53%)	
KI67 (%)			0.023
<=45	3 (33%)	27 (79%)	
>45	6 (67%)	7 (21%)	
ECOG			>0.9
0-1	10 (83%)	34 (81%)	
2	2 (17%)	8 (19%)	
LDH rate			0.2
>N	7 (58%)	13 (31%)	
N	5 (42%)	29 (69%)	
Leukocytes count			0.2
<=5G/L	6 (50%)	32 (76%)	
>5G/L	6 (50%)	10 (24%)	
Albumin			0.5
<3.6g/dL	4 (36%)	8 (21%)	
>=3.6g/dL	7 (64%)	31 (79%)	
Ann Arbor stage			0.9
II	0 (0%)	1 (2.4%)	
III	1 (8.3%)	3 (7.1%)	
IV	11 (92%)	38 (90%)	
Bulky tumor			0.9
no	8 (89%)	33 (97%)	
yes	1 (11%)	1 (2.9%)	
Extranodal involvement			>0.9
no	1 (8.3%)	4 (9.5%)	
yes	11 (92%)	38 (90%)	
Bone Marrow involvement			0.5
no	2 (17%)	13 (32%)	
yes	10 (83%)	27 (68%)	



Supplementary Figure 1 : Violin plots of percentages of positive cell for p53 by immunohistochemistry. Dashed line, positivity threshold of 50 percents.