

# Catheter-related thrombosis in stem cell recipients: comparison of different types of catheter

Slawomir Milczarek,<sup>1,2</sup> Piotr Kulig,<sup>1,2</sup> Oliwia Piotrowska,<sup>1</sup> Alina  
Zuchmanska,<sup>1</sup> Anna Bielikowicz<sup>1</sup> and Bogusław Machalinski<sup>1,2</sup>

Correspondence:

S. MAILCZAREK - slawomir.milczarek@pum.edu.pl

<sup>1</sup>Department of Hematology and Transplantology, Pomeranian  
Medical University and <sup>2</sup> Department of General Pathology,  
Pomeranian Medical University, Szczecin, Poland

<https://doi.org/10.3324/haematol.2023.283924>

**Supplementary data:**

	<b>Peripheral catheter; N/mean(median); IQR</b>	<b>Central catheter; N/mean(median); IQR</b>	<b>p- value</b>
<b>Catheter type</b>			
<i>PICC</i>	20		
<i>MLC</i>	15		
<i>CICC</i>		33	
Age	55.86(57); 42.50-66.00	59.47(61); 55.50-65.25	0.5
Sex F/M	16/19	16/17	1
<b>Diagnosis</b>			
<i>AML</i>	5		
<i>GCT</i>	8		
<i>MM</i>	21	33	
<i>WM</i>	1		
<b>Treatment regimen</b>			
<i>BuMel</i>		1	
<i>MEL 140</i>	4	2	
<i>MEL 200</i>	16	30	
<i>CE</i>	8		
<i>FluBu3</i>	1		
<i>FluBu4</i>	2		
<i>FluTBI2</i>	2		
<i>BeEAM</i>	1		
<i>BeTMI</i>	1		
Cryopreservation Yes/No	33/2	33	
<b>Type of HSCT</b>			
<i>auto-HSCT</i>	29	33	<b>0.011</b>
<i>alloHSCT</i>	6	0	
Length of stay	23.61(22.00); 21.00-26.00	21.72(21.00); 18.00-23.25	0.15

<b>Catheter size</b>			
Fr4	11 (MLC)	0	
Fr5	21	0	
Fr6	3	0	
Fr7	0	33	
<b>Number of lumens</b>			
1	5	0	
2	27	0	
3	3	33	
<b>Insertion site</b>			
Right subclavian vein	0	28	
Left subclavian vein	0	1	
Right jugular vein	0	4	
Left basilic vein	22	0	
Right basilic vein	11	0	
Right brachial vein	2	0	

**Table S1. Patients and catheter characteristics.** PCM—plasma cell myeloma, WM—Waldenström macroglobulinemia, AML—acute myeloid leukemia, GCT—germ cell tumor, Allo-HSCT—allogeneic stem cell transplantation, Auto-HSCT—autologous stem cell transplantation, MEL—melphalan, CE—carboplatin/etoposide, FluBu4—fludarabine/busulfan, BuMel—busulfan/melphalan, Fr—French gauge (1 Fr = 1/3 mm). P value was calculated where applicable. N – number, IQR – interquartile range. Statistically significant results are shown in bold.

	Estimate	OR	Standard error	Z value	p
Intercept	9.56882	14311.45	4.12663	2.319	<b>0.02</b>
Catheter length	-0.06843	0.93	0.04261	-1.606	0.11
Catheter diameter	-4.42005	0.012	1.20493	-3.668	<b>&lt; 0.001</b>
Length of stay	0.01965	1.02	0.08752	0.225	0.82
Type of HSCT	-1.70001	0.18	1.7661	-0.963	0.34

**Table S2. Multiple logistic regression model for a catheter dysfunction.** P value < 0.05 was considered statistically significant. Statistically significant results are shown in bold. OR – odds ratio.