High annualized bleeding rates in pediatric patients with inherited platelet function disorders

Authors


Shelly Saini,¹ Song Zhang,² Sean Yates,³ Jessica Garcia,^{1,4} Ruchika Sharma,^{1,4} Joseph Formella,¹ Ravindra Sarode³ and Ayesha Zia^{1,4}


¹Department of Pediatrics; ²O'Donnell School of Public Health; ³Department of Pathology and ⁴Division of Hematology/Oncology, The University of Texas Southwestern Medical Center, Dallas, TX, USA

Correspondence:

A. ZIA - Ayesha.zia@utsouthwestern.edu


https://doi.org/10.3324/haematol.2024.284996






Supplemental Figure 1. Frequency of different types of (A) hemorrhagic events and (B) treatments within each study group. Controls had a bleeding phenotype but no identifiable bleeding disorder on comprehensive laboratory evaluation.

Abbreviations: VWD, von Willebrand Disease; IPFD, inherited platelet function disorder; GI, gastroenterological; CNS, central nervous system



Supplemental Figure 2. ROC curves for prediction of bleeding outcomes. ROC curves for the prediction of (A) bleeding events, (B) bleeds requiring treatment, and (C) severe bleeding events in IPFD and VWD subjects with a bleeding score (BS) above the indicated threshold.

Abbreviations: VWD, von Willebrand Disease; IPFD, inherited platelet function disorder; AUC, area under the curve



Supplemental Figure 3. Odds ratio for likelihood of suffering bleeding outcomes. Likelihood to suffer (A) clinically significant bleeding events, (B) bleeds requiring treatment, and (C) severe bleeds (odds ratio and 95% confidence interval) for subjects with a bleeding score above the identified cutoff.

Abbreviations: VWD, von Willebrand Disease; IPFD, inherited platelet function disorder