Outcomes after chimeric antigen receptor T-cell therapy across large B-cell lymphoma subtypes

Authors

Christianne Bourlon,¹* Claire Roddie,²,³* Tobias Menne,⁴ Jane Norman,⁵ Maeve O'Reilly,² Adam Gibb,⁶ Caroline Besley,⁶ Sridhar Chaganti,⁶ Carlos Gonzalez Arias,⁶ Ceri Jones,¹⁰ Abdalla Dikair,¹¹ Sharon Allen,¹² Frances Seymour,¹³ Wendy Osborne,⁴ Amrith Mathew,⁶ William Townsend,² Piers E. M. Patten,¹¹⁴ Eleni Thoulouli,⁵ Ahmed Abdulgawad,¹⁵ Sanne Lugthart,⁶ Robin Sanderson,¹ Amy A. Kirkwood¹⁶ and Andrea Kuhnl¹

¹Department of Hematology, King's College Hospital, London; ²Department of Hematology, University College London Hospitals, London; ³UCL Cancer Institute, University College London, London; ⁴Department of Hematology, Freeman Hospital, Newcastle University, Newcastle; ⁵Department of Hematology, Manchester Royal Infirmary, Manchester; ⁶Department of Medical Oncology, The Christie Hospital, Manchester; ⁷Department of Hematology, University Hospitals Bristol and Weston, Bristol; ⁸Department of Hematology, Queen Elizabeth Hospital, Birmingham; ⁹Department of Hematology, Royal Marsden Hospital, London; ¹⁰Department of Hematology, University Hospital of Wales, Cardiff; ¹¹Department of Hematology, Queen Elizabeth Hospital, Glasgow; ¹²Department of Hematology, Cambridge University Hospitals, Cambridge; ¹³Department of Hematology, Leeds Teaching Hospitals, Leeds; ¹⁴Comprehensive Cancer Center, King's College London, London; ¹⁵Department of Hematology, The Christie Hospital, Manchester and ¹⁶CR UK & UCL Cancer Trials Center, UCL Cancer Institute, UCL, London, UK

*CB and CR contributed equally as first authors.

Correspondence:

A. KUHNL - andrea.kuhnl@nhs.net

https://doi.org/10.3324/haematol.2024.285010

Data Supplement

Outcomes after chimeric antigen receptor T-cell therapy across large B-cell lymphoma subtypes

Table S1. CAR T toxicity across subgroups

Best response	De novo DLBCL N=420	PMBCL N=26	t-FL N=130	t-NFL N=38	p-value ¹	t-MZL N=17	RS N=13	t-other N=8
Grade 3+ CRS	20/420 (4.8%)	3/26 (11.5%)	7/130 (5.4%)	2/38 (5.3%)	0.43	2/17 (11.8%)	0/13 (0%)	0/8 (0%)
Any grade CRS ²	272/319 (85.3%)	22/26 (84.6%)	75/98 (76.5%)	19/26 (73.1%)	0.11	11/13 (84.6%)	4/6 (66.7%)	4/7 (57.1%)
Grade 3+ ICANS	66/420 (15.7%)	5/26 (19.2%)	21/130 (16.2%)	3/38 (7.9%)	0.55	1/17 (5.9%)	2/13 (15.4%)	0/8 (0%)
Any grade ICANS ²	132/319 (41.4%)	9/26 (34.6%)	35/98 (35.7%)	5/26 (19.2%)	0.13	2/13 (15.4%)	3/6 (50.0%)	0/7 (0%)
Steroids used	119/274 (43.4%)	9/23 (39.1%)	36/77 (46.8%)	3/19 (15.8%)	0.088	3/13 (23.1%)	0/1 (0%)	0/5 (0%)
Tocilizumab used	199/274 (72.6%)	16/23 (69.6%)	53/78 (68.0%)	11/19 (57.9%)	0.48	7/13 (53.9%)	1/1 (100%)	3/5 (60.0%)
ICU admission	69/244 (28.3%)	7/23 (30.4%)	16/73 (21.9%)	3/18 (16.7%)	0.55	3/13 (23.1%)	0/1 (0%)	0/4 (0%)
Grade 3+ thrombocytopenia 1 month ³	57/146 (39.0%)	6/17 (35.3%)	26/54 (48.2%)	5/10 (50.0%)	0.59	4/6 (66.8%)	-	1/4 (25.0%)
Grade 3+ neutropenia 1 month ³	68/145 (46.9%)	7/16 (43.8%)	26/53 (49.1%)	5/9 (55.6%)	0.94	3/5 (60.0%)	-	2/4 (50.0%)
Grade 3+ thrombocytopenia 3 months ³	9/81 (11.1%)	2/11 (18.2%)	6/37 (16.2%)	2/5 (40.0%)	0.23	1/3 (33.3)	-	1/2 (50.0%)
Grade 3+ neutropenia 3 months ³	15/81 (18.5%)	2/11 (18.2%)	7/37 (18.9%)	2/5 (40.0%)	0.63	1/3 (33.3)	-	1/2 (50.0%)

¹Fisher's exact test comparing rates in de novo DLBCL/PMBCL/t-FL and t-NFL groups. ²Any grade: chi-squared for trend 0.13 (CRS), 0.036 (ICANS: lower grades in t-NFL patients, p=0.30 if these patients are excluded). ³Thrombocytopenia and Neutropenia rates exclude patients who have relapsed by each timepoint.