

Reference interval of free light chains ratio in patients with end-stage renal disease on chronic hemodialysis

Authors

Camila Peña,^{1,2*} Ricardo Valjalo,^{3*} Ramón Pérez,⁴ Marco Álvarez,⁵ Pablo Bustamante,⁵ Esteban Forray,⁵ Viviana Balboa⁴ and Alexis Bondí⁴

¹Hematology Section, Hospital del Salvador; ²Centro de Investigación Clínica Avanzada (CICA-Oriente); ³Nephrology Section, Hospital del Salvador; ⁴Immunology Laboratory, Hospital del Salvador and ⁵Internal Medicine Section, Hospital del Salvador, Santiago de Chile, Chile

**CP and RV contributed equally as first authors.*

Correspondence:

C PEÑA - cpena@hsalvador.cl


<https://doi.org/10.3324/haematol.2024.285093>

Supplementary Table 1. Characteristics of patients with incidentally detected monoclonal gammopathy.

Case	Sex	Age	Diagnosis	Paraprotein	M protein quantif on SPEP	Kappa/Lambda (mg/L)	FLCr
1	M	85	Multiple Myeloma	IgA Lambda	2.5 g/dL	53.2/552.3	0.09
2	M	52	MGUS (MGRS?)	IgG Lambda	1.7 g/dL	166.4/242.6	0.69
3	M	65	MGUS (MGRS?)	IgG Kappa (By ISUB)	0 g/dL	590.2/149.2	3.96
4	F	75	MGUS (MGRS?)	Kappa	0 g/dL	326.8/82.9	3.94
5	M	95	Multiple Myeloma	IgG Kappa, IgA Kappa	1.4 g/dL, 0.4 g/dL	1224.2/73.9	16.56

MGUS=monoclonal gammopathy of undetermined significance. MGRS=monoclonal gammopathy of renal significance. ISUB= immunosubstraction. FLCr=Serum Free Light Chain Ratio.

Supplementary Figure 1. Pre-dialysis Kappa and Lambda serum free light chains (sFLC) concentrations in the 137 patients with HD-dependent end stage renal disease.



Supplementary Figure 2. Kappa and Lambda concentrations in patients with and without residual kidney function (RKF).

