Comparison of EasyM, a clonotypic mass spectrometry assay, and EuroFlow minimal residual disease assessment in multiple myeloma

Authors

Jessie Zhao,^{1,2} Tiffany Khong,^{1,2} Malgorzata Gorniak,³ Abir Khaled,⁴ Zac McDonald,⁴ John Reynolds,^{1,2} Sridurga Mithraprabhu,^{1,2} Nicholas Bingham,^{1,2} Suehli Lim,^{1,2} Daniel Wong,^{1,2} Anna Johnston,⁵ Olga Motorna,⁶ Nicholas Murphy,⁵ Hang Quach,⁷ Liqiang Yang⁴ and Andrew Spencer^{1,2}

¹Department of Malignant Hematology & Stem Cell Transplantation, The Alfred Hospital, Melbourne, Victoria, Australia; ²Australian Center for Blood Diseases, Monash University, Melbourne, Victoria, Australia; ³Alfred Pathology, The Alfred Hospital, Melbourne, Victoria, Australia; ⁴Rapid Novor, Kitchener, Ontario, Canada; ⁵Department of Hematology, Royal Hobart Hospital, Hobart, Tasmania, Australia; ⁶Department of Hematology, Eastern Health, Melbourne, Victoria, Australia and

⁷Department of Hematology, St. Vincent's Hospital Melbourne, Melbourne, Victoria, Australia

Correspondence:

A. SPENCER - andrew.spencer@monash.edu


https://doi.org/10.3324/haematol.2024.285933

Received: May 28, 2024. Accepted: October 29, 2024. Early view: November 7, 2024.

©2025 Ferrata Storti Foundation

Published under a CC BY-NC license

Supplementary Figure 1. **MM19 and MM21 trial schema**. ASCT, autologous stem cell transplantation; DRd, daratumumab-lenalidomide-dexamethasone; ITd, ixazomib-thalidomide-dexamethasone; MM, multiple myeloma; MRD, minimal residual disease; MS, mass spectrometry; NGF, next-generation flow cytometry; PD, progressive disease; PR, partial response; R, lenalidomide; Td, thalidomide-dexamethasone.



Supplementary Figure 2. **Overview of the EasyM assay**. LC-MS/MS, liquid chromatography with tandem mass spectrometry; M-protein, monoclonal protein.

